

DEX-0113



BM

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 9/64, 15/09, 15/57, C07K 16/40, 16/18, A61K 38/48, 39/395	A1	(11) International Publication Number: WO 98/13484 (43) International Publication Date: 2 April 1998 (02.04.98)
(21) International Application Number: PCT/US97/17175 (22) International Filing Date: 25 September 1997 (25.09.97) (30) Priority Data: 08/723,938 26 September 1996 (26.09.96) US (71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BANDMAN, Olga [US/US]; 366 Anna Avenue, Mountain View, CA 94043 (US). COLEMAN, Roger [US/US]; 260 Mariposa #2, Mountain View, CA 94041 (US). (74) Agent: BILLINGS, Lucy, J.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US).		(81) Designated States: AT, AU, BR, CA, CH, CN, ES, FI, GB, IL, JP, KR, MX, NO, NZ, RU, SE, SG, US, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: TWO HUMAN CATHEPSIN PROTEINS (57) Abstract <p>The present invention provides two novel human cathepsin proteins (HCPs) and polynucleotides encoding HCPs. The invention provides for genetically engineered expression vectors and host cells comprising the nucleic acid sequence encoding HCPs. The invention also provides for the production and use of antibodies to HCPs in pharmaceutical compositions for the treatment of disease processes that include cancers, inflammation, metastasis and peptide and proenzyme processing. In addition, the invention provides for the production and use of inhibitors of HCPs in pharmaceutical compositions for the treatment of diseases. The invention also describes diagnostic assays which utilize the polynucleotide to hybridize with the transcripts encoding HCPs. The invention also provides for the use of antisense molecules in pharmaceutical compositions as therapeutics in cancers, inflammation, metastasis and peptide and proenzyme processing.</p>		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TWO HUMAN CATHEPSIN PROTEINS

TECHNICAL FIELD

The present invention relates to nucleic acid and amino acid sequences of two novel
5 human cathepsin proteins and to the use of these sequences in the diagnosis, study, prevention
and treatment of disease.

BACKGROUND ART

The cathepsin family of lysosomal protease includes the cysteine protease; cathepsins B,
H, K, L, O2, and S, and the aspartyl protease; cathepsins D, and G. The various members of this
10 endosomal protease family are differentially expressed. Some, such as cathepsin D, have a
ubiquitous tissue distribution while others, such as cathepsin L, are found only in monocytes,
macrophages, and other cells of the immune system.

The cathepsins represent the major endopeptidases in the lysosome (Huisman W, et al.
(1974) *Biochem Biophys Acta* 370:297-307) and as such participate in the degradation of
15 proteins entering the vacuolar system by endocytosis and in the turnover of cytoplasmic proteins.
The cathepsins are also active in 1) initiation of proteolytic cascades by proenzyme activation, 2)
processing of the II alpha beta heterodimer in endosomes for expression of functional MHC class
II molecules which bind antigenic peptides, and 3) processing of antigen in antigen-presenting
cells. The secreted forms of several members of this family function in tissue remodeling
20 through degradation of collagen, laminin, elastin, and other structural proteins of basement
membranes (Mizuochi T, (1994) *Immunol Lett* 43:189-193, Baldwin E T. (1993) *Proc Natl Acad*
Sci 90:6796-6800). Both cysteine and aspartyl cathepsins are used by various parasitic protozoa
for the catabolism of host cell proteins and/or facilitating host invasion (Becker M M et al. (1995)
J Biol Chem 270 24496-24501, Rosenthal P J et al. (1988) *J Clin Invest* 82:1560-1566,
25 McKerrow J H (1993) *Annu Rev Microbiol* 47:821-853).

The various cathepsin proteases differ in their gene structures and in their transcriptional
regulation. The cathepsin D gene promoter has a compound structure with features of both
housekeeping genes (high G+C content and potential transcription factor SP-1 sites) and
regulated genes (TATA sequence). RNase protection assays show that transcription is initiated at
30 five major transcription sites (transcription site-I to transcription site-V) spanning 52 base pairs.
Site-directed mutagenesis studies indicate that the TATA box is essential for initiation of
cathepsin D gene transcription at transcription site-I. This suggests that cathepsin D behaves,

depending on the conditions, as a housekeeping gene with multiple start sites or as a hormone-regulated gene that can be controlled from its TATA box (Cavaillès V (1993) *Proc Natl Acad Sci* 90:203-207). The cathepsin L gene promoter has no TATA box but includes several SP-1 sites, two AP-2 transcription regulatory element binding sites (a promoter under the control of the
5 tumor promoter and cAMP), and a cAMP response element. Experimental data confirm that the expression of cathepsin L is induced by malignant transformation, growth factors, tumor promoters, and cyclic AMP (Troen B et al. (1991) *Cell Growth Differ* 2:23-31).

Altered regulation and expression of these two different cathepsins is evident in disease states. Cathepsin D is overproduced and hypersecreted by breast cancer cells. Clinical studies
10 show a strong correlation between high concentrations of cathepsin D in the cytosol of primary tumor cells and further occurrence of metastasis. Cathepsin D is expressed at high levels in hormone independent breast cancer, is induced by estrogen in hormone dependent breast cancer, and appears to be independent of other more classical prognostic factors. In nude mice, transfection of cathepsin D cDNA into tumor cells increases their metastatic potential, suggesting
15 that overexpression of this protease may be one of the factors responsible for metastasis (Rochefort H (1992) *Acta Oncol* 31:125-30, Long B J (1996) *Cancer Lett* 99:233-238). The mechanism by which this protease might facilitate metastasis is not fully characterized, although cathepsin D has the potential to initiate a proteolytic cascade, to degrade extracellular matrix and to liberate growth factors from the matrix. In vitro studies have noted that transfected cathepsin
20 D stimulates high density cancer cell growth via an intracellular mechanism leading to a decreased secretion of growth inhibitors (Liaudet E (1995) *Cell Growth Differ* 6:1045-1052). Altered cathepsin D levels are present in biopsy specimens in prostate and bladder carcinomas and are shown to correlate with tumor grade (Ross J S (1995) *Am J Clin Pathol* 104:36-41, Dickinson A J (1995) *J Urol* 154:237-241).

25 Altered cathepsin activity and/or distribution may play a role in certain brain diseases. In Alzheimer's disease, A4 amyloid peptide, the main constituent of amyloid plaques and cerebrovascular amyloid deposits, derives from a large amyloid precursor protein (APP) by the action of endoproteases which release the amino and carboxyl termini to generate the aggregating form of A4. In the brains of Alzheimer's patients, more than 90% of the pyramidal neurons in
30 lamina V and 70% in lamina III displayed 2- to 5-fold elevated levels of cathepsin D mRNA by in situ hybridization compared with neurologically normal controls. An endogenous protease activity from diseased samples was found to be active in acidic conditions and inhibited by

pepstatin, two characteristics of cathepsin D, suggesting that a cathepsin D-like activity from human brain may be responsible for APP processing (Evin G (1995) *Biochemistry* 34:14185-14192, Cataldo A M (1995) *Neuron* 14:671-680). There is a significant increase in cathepsin D activity in biopsies from Huntington's disease (Mantle D (1995) *J Neurol Sci* 131:65-70) and
5 there are increased levels of cathepsin D mRNA in scrapie-infected mice (Diedrich J F (1991) *J Virol* 65:4759-4768).

Abnormal regulation of cathepsins is observed in several inflammatory disease states. In fibroblastoid synovial lining cells isolated from rheumatoid and other chronic inflammatory synovial tissues, the mRNA for stromelysin, vimentin, IL-4, IL-6, TIMP-1, cathepsin D,
10 gelatinase, TGF alpha, c-fms and DR beta is preferentially expressed. This modulated pattern of gene expression is correlated with the phenotype of this inflammatory condition (Ritchlin C et al. (1994) *Scand J Immunol* 40:292-298). Cathepsin L and D expression was evaluated in synovial tissues from patients with rheumatoid arthritis (RA) and osteoarthritis (OA), using in situ hybridization with digoxigenin-labeled RNA probes. Both RA and OA synovial tissue expressed
15 cathepsins L and D. The expression of the cathepsins was markedly higher in interstitial regions and in perivascular infiltrates of RA synovial tissue compared with OA specimens. The differential expression of cathepsins L and D mRNA in RA and OA synovial tissues supports the concept that these enzymes may contribute to the influx of mononuclear cells into the synovium and suggests that the adhesion of synovial cells to cartilage mediates the invasive/destructive
20 process in RA (Keyszer G M (1995) *Arthritis Rheum* 38:976-984).

The cathepsins are believed to be involved in several other diseases states. In an experimental model of human glomerular disease, the administration of a specific, irreversible inhibitor of cysteine protease (trans-epoxysuccinyl-L-leucylamido-(3-methyl)butane) significantly reduces proteinuria in rats (Baricos W H (1991) *Arch Biochem Biophys* 288:468-
25 72.) The fibroblasts from patients with mucopolipidosis II and III have a severely compromised capacity for endogenous lysosomal protein degradation that appears to result from multiple cathepsin deficiencies (Kopitz J (1993) *Biochem J* 295 (Pt 2): 577-580). The platelet aggregating cysteine protease implicated in thrombotic thrombocytopenic purpura shows the characteristics of a lysosomal cathepsin (Consonni R (1994) *Br J Haematol* 87:321-324).

30 Cathepsin D knockout mice develop normally during the first 2 weeks, stop thriving in the third week and die in a state of anorexia at day 26 +/- 1. An atrophy of the ileal mucosa observed in the third week progresses towards widespread intestinal necroses accompanied by

thromboemboli. The thymus and spleen undergo massive destruction with loss of T and B cells. The lysosomal bulk proteolysis is, however, maintained. These results suggest that the major functions of cathepsin D involve limited proteolysis of proteins regulating cell growth and/or tissue homeostasis (Saftig P, (1995) EMBO J 14:3599-3608).

5 Cathepsins have a role in processes that involve proteolysis of specific proteins and tissues in local microenvironments including inflammation, metastasis and peptide and proenzyme processing. The increased expression and differential regulation of these protease is linked to the metastatic potential of a variety of cancers and as such is of therapeutic and prognostic interest. Evidence of the involvement of cathepsins associated with protein
10 processing in diseases such as Alzheimer's disease, Huntington's disease, mucopolidosis and arthritic inflammation suggests that modulation of the cathepsins may ameliorate these disease processes. The polynucleotide sequences and proteins of the claimed invention would satisfy this need by providing the means for diagnosis, study, prevention and treatment of these diseases.

DISCLOSURE OF THE INVENTION

15 The present invention discloses two novel human cathepsin proteins, hereinafter referred to as HCP-1 and HCP-2 and, collectively, HCPs, which share features with other proteins involved in proteolysis. Accordingly, the invention features substantially purified HCP-1 and HCP-2, as shown in the amino acid sequence of SEQ ID NO:1 and SEQ ID NO:5.

 One aspect of the invention features isolated and substantially purified polynucleotides
20 which encode HCPs. In a particular aspect, the polynucleotides are the nucleotide sequences of SEQ ID NO:1 and SEQ ID NO:5. In addition, the invention features polynucleotide sequences that hybridize under stringent conditions to SEQ ID NO:1 and SEQ ID NO:5.

 The invention further relates to the nucleic acid sequence encoding HCP, oligonucleotides, peptide nucleic acids (PNA), fragments, portions or antisense molecules
25 thereof. The present invention also relates, in part, to the inclusion of the nucleic acid sequence encoding HCP in an expression vector which can be used to transform host cells.

 The present invention also relates to a method for producing HCP or a fragment thereof. It contemplates the delivery of purified, alone or in a pharmaceutically acceptable excipient, to cancerous cells or tissues. It also encompasses antibodies which bind specifically to HCP and
30 can be used to monitor testing of cathepsin-expressing tissues.

BRIEF DESCRIPTION OF DRAWINGS

Figures 1A, 1B, 1C, 1D and 1E shows the amino acid sequence (SEQ ID NO:1) and

nucleic acid sequence (SEQ ID NO:2) for HCP-1.

Figures 2A and 2B shows the amino acid alignment of HCP-1 and GI 555662 (Smith A M,(1994)Mol Biochem Parasitol 67 (1):11-19) The alignment was produced using the multisequence alignment program of DNASTar™ software (DNASTar Inc, Madison WI).

5 Figure 3 shows the hydrophobicity plot for HCP-1; the X axis reflects amino acid position, and the negative Y axis, hydrophobicity (MacDNAsis software, Hitachi Software Engineering Co Ltd, San Bruno CA).

Figure 4 shows the hydrophobicity plot for GI 555662; (MacDNAsis software).

Figures 5A, 5B, 5C and 5D shows the amino acid sequence (SEQ ID NO:3) and nucleic
10 acid sequence (SEQ ID NO:4) for HCP-2.

Figures 6A and 6B shows the amino acid alignment of HCP-2 and GI 181180 (Faust P L (1985)Proc Natl Acad Sci 82 (15)4910-4914). (DNASTar Inc, Madison WI).

Figure 7 shows the isoelectric plot for HCP-2 (MacDNAsis software).

Figure 8 shows the isoelectric plot for GI 181180 (MacDNAsis software).

15 **MODES FOR CARRYING OUT THE INVENTION**

Definitions

"Nucleic acid sequence" as used herein refers to an oligonucleotide, nucleotide or polynucleotide, and fragments or portions thereof, and to DNA or RNA of genomic or synthetic origin which may be single- or double-stranded, and represent the sense or antisense strand.

20 Similarly, amino acid sequence as used herein refers to peptide or protein sequence.

"Consensus" as used herein may refer to a nucleic acid sequence 1) which has been resequenced to resolve uncalled bases, 2) which has been extended using XL-PCR (Perkin Elmer) in the 5' or the 3' direction and resequenced, 3) which has been assembled from the overlapping sequences of more than one Incyte clone GCG Fragment Assembly System, (GCG,
25 Madison WI), or 4) which has been both extended and assembled.

"Peptide nucleic acid" as used herein refers to a molecule which comprises an oligomer to which an amino acid residue, such as lysine, and an amino group have been added. These small molecules, also designated anti-gene agents, stop transcript elongation by binding to their complementary (template) strand of nucleic acid (Nielsen PE et al (1993) Anticancer Drug Des
30 8:53-63).

A "deletion" is defined as a change in either nucleotide or amino acid sequence in which one or more nucleotides or amino acid residues, respectively, are absent.

An "insertion" or "addition" is that change in a nucleotide or amino acid sequence which has resulted in the addition of one or more nucleotides or amino acid residues, respectively, as compared to the naturally occurring HCP.

A "substitution" results from the replacement of one or more nucleotides or amino acids
5 by different nucleotides or amino acids, respectively.

As used herein, HCP refers to the amino acid sequence of substantially purified HCP obtained from any species, particularly mammalian, including bovine, ovine, porcine, murine, equine, and preferably human, from any source whether natural, synthetic, semi-synthetic or recombinant.

10 A "variant" of HCP is defined as an amino acid sequence differs by one or more amino acids. The variant may have "conservative" changes, wherein a substituted amino acid has similar structural or chemical properties, eg, replacement of leucine with isoleucine. More rarely, a variant may have "nonconservative" changes, eg, replacement of a glycine with a tryptophan. Similar minor variations may also include amino acid deletions or insertions, or both. Guidance
15 in determining which and how many amino acid residues may be substituted, inserted or deleted without abolishing biological or immunological activity may be found using computer programs well known in the art, for example, DNASTar software.

The term "biologically active" refers to a HCP having structural, regulatory or biochemical functions of naturally occurring HCP. Likewise, "immunologically active" defines
20 the capability of the natural, recombinant or synthetic HCP, or any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The term "derivative" as used herein refers to the chemical modification of nucleic acids encoding HCP or the encoded HCP. Illustrative of such modifications would be replacement of
25 hydrogen by an alkyl, acyl, or amino group. A nucleic acid derivative would encode a polypeptide which retains essential biological characteristics of natural HCP.

As used herein, the term "substantially purified" refers to molecules, either nucleic or amino acid sequences, that are removed from their natural environment, isolated or separated, and are at least 60% free, preferably 75% free, and most preferably 90% free from other
30 components with which they are naturally associated.

The term "hybridization" as used herein shall include "any process by which a strand of nucleic acid joins with a complementary strand through base pairing" (Coombs J (1994)

Dictionary of Biotechnology, Stockton Press, New York NY). Amplification is defined as the production of additional copies of a nucleic acid sequence and is generally carried out using polymerase chain reaction technologies well known in the art (Dieffenbach CW and GS Dveksler (1995) PCR Primer, a Laboratory Manual, Cold Spring Harbor Press, Plainview NY).

- 5 "Stringency" typically occurs in a range from about $T_m - 5^\circ\text{C}$ (5°C below the T_m of the probe) to about 20°C to 25°C below T_m . As will be understood by those of skill in the art, a stringency hybridization can be used to identify or detect identical polynucleotide sequences or to identify or detect similar or related polynucleotide sequences.

Preferred Embodiments

- 10 The consensus nucleotide sequence for HCP-1 (SEQ ID NO:1), disclosed herein, encodes a novel human cathepsin protein of 392 amino acid residues. The consensus sequence is based on the extension and assembly of the following Incyte clones: 152940 (FIBRAGT02), 398290 (PITUNOT02), 723324 (SYNOOAT01), 787555 (PROSNOT05), and 937823 (CERVNOT01). HCP-1 has 392 amino acid residues including seven conserved cysteines. In addition, HCP-1 has
 15 49% identity to *Schistosoma mansoni* puerto rican preprocathepsin L (GI 555662, Figure 2A and 2B) as well as similarities in hydrophobicity (Figures 4, 5A, 5B, 5C and 5D). The amino acid alignments of the consensus sequence and GI 555662 are shown in Figures 2A and 2B. Using the numbers for the consensus sequence at the side of the figure as a reference, the following cysteine residues are conserved: C₂₀₀, C₂₀₃, C₂₃₄, C₂₄₁, C₂₇₄, C₃₃₂, and C₃₈₀. The hydrophobicity plot for the
 20 consensus sequence aligns with that for GI 555662 further suggesting similar functions as a cysteine protease.

- The Incyte clone number 312099, HCP-2, encodes a novel human cathepsin protein of 395 amino acid residues (Figures 5A, 5B, 5C and 5D). The clone sequence is based on the extension of Incyte clone number 312099 (LUNGNOT02). HCP-2 has 395 amino acid residues
 25 including nine conserved aspartic acids, D₉₆, D₁₅₃, D₁₈₄, D₂₁₃, D₂₂₄, D₂₃₉, D₂₈₃, D₃₅₀ and D₃₆₁ and four conserved cysteines, C₁₀₉, C₁₁₆, C₂₇₄, and C₃₁₇. HCP-2 has 43% identity with human preprocathepsin D (GI 181180, Fig 6) and a similar isoelectric point (Figures 7 and 8).

The HCP Coding Sequences

- The nucleic acid and deduced amino acid sequences of the HCP-1 is shown in Figures
 30 1A, 1B, 1C, 1D and 1E. In accordance with the invention, any nucleic acid sequence which encodes the HCP can be used to generate recombinant molecules which express HCP. In a specific embodiment described herein, a partial sequence encoding HCP-1 was first isolated as

Incyte Clone 152940 from an ataxia telangiectasia fibroblast cDNA library (FIBRAGT02).

The nucleic acid and deduced amino acid sequences of the HCP-2 is shown in Figures 5A, 5B, 5C and 5D. In accordance with the invention, any nucleic acid sequence which encodes the HCP can be used to generate recombinant molecules which express HCP. In a specific
5 embodiment described herein, a partial sequence encoding HCP-2 was first isolated as Incyte Clone 312099 from a cDNA library (LUNGNOT02). It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of HCP-encoding nucleotide sequences, some bearing minimal homology to the nucleotide sequences of any known and naturally occurring gene may be produced. The invention contemplates each and
10 every possible variation of nucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the nucleotide sequence encoding naturally occurring HCP, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode HCP and their variants are preferably
15 capable of hybridizing to the nucleotide sequence of the naturally occurring sequences under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding HCP or their derivatives possessing a substantially different codon usage. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic expression host in accordance with the frequency with which
20 particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding HCP and their derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

A DNA sequence, or portions thereof, encoding HCP and their derivatives may be
25 produced entirely by synthetic chemistry, after which the synthetic gene may be inserted into any of the many available DNA vectors and cell systems using reagents that are generally available. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding HCP or any portion thereof.

Also included within the scope of the present invention are polynucleotide sequences that
30 are capable of hybridizing to the nucleotide sequences of SEQ ID NO:2 and SEQ ID NO:4 under various conditions of stringency. Hybridization conditions are based on the melting temperature (T_m) of the nucleic acid binding complex or probe, as taught in Berger and Kimmel (1987,

Guide to Molecular Cloning Techniques, Methods in Enzymology, Vol 152, Academic Press, San Diego CA) incorporated herein by reference, and may be used at a defined "stringency".

Altered nucleic acid sequences encoding HCP which may be used in accordance with the invention include deletions, insertions or substitutions of different nucleotides resulting in a polynucleotide that encodes the same or a functionally equivalent HCP. The protein may also show deletions, insertions or substitutions of amino acid residues which produce a silent change and result in functionally equivalent HCP. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues as long as the biological activity of HCP is retained. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine, valine; glycine, alanine; asparagine, glutamine; serine, threonine, phenylalanine, and tyrosine.

Included within the scope of the present invention are alleles encoding HCP. As used herein, an "allele" or "allelic sequence" is an alternative form of the nucleic acid sequence encoding HCP. Alleles result from a mutation, ie, a change in the nucleic acid sequence, and generally produce altered mRNAs or polypeptides whose structure or function may or may not be altered. Any given gene may have none, one or many allelic forms. Common mutational changes which give rise to alleles are generally ascribed to natural deletions, additions or substitutions of amino acids. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

Methods for DNA sequencing may be used which are well known in the art and employ such enzymes as the Klenow fragment of DNA polymerase I, Sequenase® (US Biochemical Corp, Cleveland OH), Taq polymerase (Perkin Elmer, Norwalk CT), thermostable T7 polymerase (Amersham, Chicago IL), or combinations of recombinant polymerases and proofreading exonucleases such as the ELONGASE Amplification System marketed by Gibco BRL (Gaithersburg MD). Preferably, the process is automated with machines such as the Hamilton Micro Lab 2200 (Hamilton, Reno NV), Peltier Thermal Cycler (PTC200; MJ Research, Watertown MA) and the ABI 377 DNA sequencers (Perkin Elmer).

30 Extending the Polynucleotide Sequence

The polynucleotide sequences encoding HCP may be extended utilizing partial nucleotide sequence and various methods known in the art to detect upstream sequences such as promoters

and regulatory elements. For example, the method of Gobinda et al (1993; PCR Methods Applic 2:318-22) involves "restriction-site" polymerase chain reaction (PCR) as a direct method which uses universal primers to retrieve unknown sequence adjacent to a known locus. First, genomic DNA is amplified in the presence of primer to a linker sequence and a primer specific to the
5 known region. The amplified sequences are subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

Inverse PCR can be used to amplify or extend sequences using divergent primers based on a known region (Triglia T et al (1988) Nucleic Acids Res 16:8186). The primers may be
10 designed using OLIGO® 4.06 Primer Analysis Software (1992; National Biosciences Inc, Plymouth MN), or another appropriate program, to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68°-72° C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR
15 template.

Capture PCR (Lagerstrom M et al (1991) PCR Methods Applic 1:111-19) may also be used as a method for PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA. Capture PCR also requires multiple restriction enzyme digestions and ligations to place an engineered double-stranded sequence into an
20 unknown portion of the DNA molecule before PCR.

Another method which may be used to retrieve unknown sequence is walking PCR (Parker JD et al (1991) Nucleic Acids Res 19:3055-60), and which involves targeted gene walking. Alternatively, PCR, nested primers, PromoterFinder™ (Clontech, Palo Alto CA) and PromoterFinder libraries can be used to walk in genomic DNA. This process avoids the need to
25 screen libraries and is useful in finding intron/exon junctions.

Preferred libraries for screening for full length cDNAs are those which have been size-selected to include larger cDNAs. Also, random primed libraries are preferred in that they will contain more sequences which contain the 5' and upstream regions of genes. A randomly primed library may be particularly useful if an oligo d(T) library does not yield a full-length
30 cDNA. Genomic libraries are useful for extension into the 5' nontranslated regulatory region.

Capillary electrophoresis may be used to analyze either the size or confirm the nucleotide sequence in sequencing or PCR products. Systems for rapid sequencing are available from

Perkin Elmer, Beckman Instruments (Fullerton CA), and other companies. Capillary sequencing may employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity is converted to electrical signal using
5 appropriate software (eg. Genotyper™ and Sequence Navigator™ from Perkin Elmer) and the entire process from loading of samples to computer analysis and electronic data display is computer controlled. Capillary electrophoresis is particularly suited to the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample. The reproducible sequencing of up to 350 bp of M13 phage DNA in 30 min has been reported
10 (Ruiz-Martinez MC et al (1993) Anal Chem 65:2851-8).

Expression of the Nucleotide Sequence

In accordance with the present invention, polynucleotide sequences which encode HCP, fragments of the polypeptides, fusion proteins or functional equivalents thereof may be used in recombinant DNA molecules that direct the expression of HCP in appropriate host cells. Due to
15 the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence, may be used to clone and express HCP. As will be understood by those of skill in the art, it may be advantageous to produce HCP-encoding nucleotide sequences possessing non-naturally occurring codons. Codons preferred by a particular prokaryotic or eukaryotic host (Murray E et al (1989) Nuc Acids Res
20 17:477-508) can be selected, for example, to increase the rate of HCP expression or to produce recombinant RNA transcripts having desirable properties, such as a longer half-life, than transcripts produced from naturally occurring sequence.

The nucleotide sequences of the present invention can be engineered in order to alter HCP-encoding sequences for a variety of reasons, including but not limited to, alterations which
25 modify the cloning, processing and/or expression of the gene product. For example, mutations may be introduced using techniques which are well known in the art, eg, site-directed mutagenesis to insert new restriction sites, to alter glycosylation patterns, to change codon preference, to produce splice variants, etc.

In another embodiment of the invention, a natural, modified or recombinant HCP-
30 encoding sequences may be ligated to heterologous sequences to encode fusion proteins. For example, for screening of peptide libraries for inhibitors of HCP activity, it may be useful to encode a chimeric HCP protein that is recognized by a commercially available antibody. A

fusion protein may also be engineered to contain a cleavage site located between an HCP and the heterologous protein sequence, so that the HCP may be cleaved and substantially purified away from the heterologous moiety.

In an alternate embodiment of the invention, the sequences encoding HCP may be synthesized, whole or in part, using chemical methods well known in the art (see Caruthers MH et al (1980) Nuc Acids Res Symp Ser 215-23, Horn T et al (1980) Nuc Acids Res Symp Ser 225-32, etc). Alternatively, the protein itself may be produced using chemical methods to synthesize a HCP amino acid sequence, whole or in part. For example, peptide synthesis can be performed using various solid-phase techniques (Roberge JY et al (1995) Science 269:202-204) and automated synthesis may be achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin Elmer) in accordance with the instructions provided by the manufacturer.

The newly synthesized peptide can be substantially purified by preparative high performance liquid chromatography (eg, Creighton (1983) Proteins, Structures and Molecular Principles, WH Freeman and Co, New York NY). The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing (eg, the Edman degradation procedure; Creighton, supra). Additionally the amino acid sequences of HCP, or any part thereof, may be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

Expression Systems

In order to express a biologically active HCP, the nucleotide sequence encoding HCP or its functional equivalent, is inserted into an appropriate expression vector, ie, a vector which contains the necessary elements for the transcription and translation of the inserted coding sequence.

Methods which are well known to those skilled in the art can be used to construct expression vectors containing HCP-encoding sequence and appropriate transcriptional or translational controls. These methods include in vitro recombinant DNA techniques, synthetic techniques and in vivo recombination or genetic recombination. Such techniques are described in Sambrook et al (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY and Ausubel FM et al (1989) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY.

A variety of expression vector/host systems may be utilized to contain and express HCP-encoding sequences. These include but are not limited to microorganisms such as bacteria

transformed with recombinant bacteriophage, plasmid or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (eg, baculovirus); plant cell systems transfected with virus expression vectors (eg, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with bacterial
5 expression vectors (eg, Ti or pBR322 plasmid); or animal cell systems.

The "control elements" or "regulatory sequences" of these systems vary in their strength and specificities and are those nontranslated regions of the vector, enhancers, promoters, and 3' untranslated regions, which interact with host cellular proteins to carry out transcription and translation. Depending on the vector system and host utilized, any number of suitable
10 transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the Bluescript® phagemid (Stratagene, La Jolla CA) or pSport1 (Gibco BRL) and ptrp-lac hybrids and the like may be used. The baculovirus polyhedrin promoter may be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (eg, heat
15 shock, RUBISCO; and storage protein genes) or from plant viruses (eg, viral promoters or leader sequences) may be cloned into the vector. In mammalian cell systems, promoters from the mammalian genes or from mammalian viruses are most appropriate. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding HCP, vectors based on SV40 or EBV may be used with an appropriate selectable marker.

20 In bacterial systems, a number of expression vectors may be selected depending upon the use intended for HCP. For example, when large quantities of HCP are needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified may be desirable. Such vectors include, but are not limited to, the multifunctional *E. coli* cloning and expression vectors such as Bluescript® (Stratagene), in which the sequence
25 encoding HCP may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke & Schuster (1989) J Biol Chem 264:5503-5509); and the like. pGEX vectors (Promega, Madison WI) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be
30 purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems are designed to include heparin, thrombin or factor XA protease cleavage sites so that the cloned polypeptide of interest can be

released from the GST moiety at will.

In the yeast, Saccharomyces cerevisiae, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase and PGH may be used. For reviews, see Ausubel et al (supra) and Grant et al (1987) *Methods in Enzymology* 153:516-544.

5 In cases where plant expression vectors are used, the expression of a sequence encoding HCP may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV (Brisson et al (1984) *Nature* 310:511-514) may be used alone or in combination with the omega leader sequence from TMV (Takamatsu et al (1987) *EMBO J* 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO (Coruzzi et al
10 (1984) *EMBO J* 3:1671-1680; Broglie et al (1984) *Science* 224:838-843); or heat shock promoters (Winter J and Sinibaldi RM (1991) *Results Probl Cell Differ* 17:85-105) may be used. These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. For reviews of such techniques, see Hobbs S or Murry LE in McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill New York NY, pp
15 191-196 or Weissbach and Weissbach (1988) Methods for Plant Molecular Biology, Academic Press, New York NY, pp 421-463.

An alternative expression system which may be used to express HCP is an insect system. In one such system, Autographa californica nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in Spodoptera frugiperda cells or in Trichoplusia larvae. The
20 sequence encoding HCP may be cloned into a nonessential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of the sequence encoding HCP will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein coat. The recombinant viruses are then used to infect S. frugiperda cells or Trichoplusia larvae in which HCP is expressed (Smith et al (1983) *J Virol* 46:584;
25 Engelhard EK et al (1994) *Proc Nat Acad Sci* 91:3224-7).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, a sequence encoding HCP may be ligated into an adenovirus transcription/ translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a nonessential E1 or E3 region of the viral genome will
30 result in a viable virus capable of expressing in infected host cells (Logan and Shenk (1984) *Proc Natl Acad Sci* 81:3655-59). In addition, transcription enhancers, such as the rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

Specific initiation signals may also be required for efficient translation of a sequence encoding HCP. These signals include the ATG initiation codon and adjacent sequences. In cases where the sequence encoding HCP, its initiation codon and upstream sequences are inserted into the most appropriate expression vector, no additional translational control signals may be needed.

5 However, in cases where only coding sequence, or a portion thereof, is inserted, exogenous translational control signals including the ATG initiation codon must be provided. Furthermore, the initiation codon must be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons can be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers

10 appropriate to the cell system in use (Scharf D et al (1994) Results Probl Cell Differ 20:125-62; Bittner et al (1987) Methods in Enzymol 153:516-544).

In addition, a host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation,

15 glycosylation, phosphorylation, lipidation and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be important for correct insertion, folding and/or function. Different host cells such as CHO, HeLa, MDCK, 293, WI38, etc have specific cellular machinery and characteristic mechanisms for such post-translational activities and may be chosen to ensure the correct modification and processing of the introduced, foreign protein.

20 For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express HCP may be transformed using expression vectors which contain viral origins of replication or endogenous expression elements and a selectable marker gene. Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose

25 of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler M et al (1977)

30 Cell 11:223-32) and adenine phosphoribosyltransferase (Lowy I et al (1980) Cell 22:817-23) genes which can be employed in tk- or aprt- cells, respectively. Also, antimetabolite, antibiotic or herbicide resistance can be used as the basis for selection; for example, dhfr which confers

resistance to methotrexate (Wigler M et al (1980) Proc Natl Acad Sci 77:3567-70); npt, which confers resistance to the aminoglycosides neomycin and G-418 (Colbere-Garapin F et al (1981) J Mol Biol 150:1-14) and als or pat, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, supra). Additional selectable genes have been described, 5 for example, trpB, which allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine (Hartman SC and RC Mulligan (1988) Proc Natl Acad Sci 85:8047-51). Recently, the use of visible markers has gained popularity with such markers as anthocyanins, β glucuronidase and its substrate, GUS, and luciferase and its substrate, luciferin, being widely used not only to identify transformants, but also to quantify the amount of 10 transient or stable protein expression attributable to a specific vector system (Rhodes CA et al (1995) Methods Mol Biol 55:121-131).

Identification of Transformants Containing the Polynucleotide Sequence

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, its presence and expression should be confirmed. For example, if the 15 sequence encoding HCP is inserted within a marker gene sequence, recombinant cells containing the sequence encoding HCP can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with the sequence encoding HCP under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem sequence as well.

20 Alternatively, host cells which contain the sequence encoding HCP and expressing HCP may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridization and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of the nucleic acid or protein.

25 The presence of the polynucleotide sequence encoding HCP can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes, portions or fragments of the sequence encoding HCP. Nucleic acid amplification based assays involve the use of oligonucleotides or oligomers based on the nucleic acid sequence to detect transformants containing DNA or RNA encoding HCP. As used herein "oligonucleotides" or "oligomers" refer 30 to a nucleic acid sequence of at least about 10 nucleotides and as many as about 60 nucleotides, preferably about 15 to 30 nucleotides, and more preferably about 20-25 nucleotides which can be used as a probe or primer.

A variety of protocols for detecting and measuring the expression of HCP, using either polyclonal or monoclonal antibodies specific for the protein are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and fluorescent activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal
5 antibodies reactive to two non-interfering epitopes on HCP is preferred, but a competitive binding assay may be employed. These and other assays are described, among other places, in Hampton R et al (1990, Serological Methods, a Laboratory Manual, APS Press, St Paul MN) and Maddox DE et al (1983, J Exp Med 158:1211).

A wide variety of labels and conjugation techniques are known by those skilled in the art
10 and can be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting related sequences include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the HCP-encoding sequence, or any portion of it, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be
15 used to synthesize RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3 or SP6 and labeled nucleotides.

A number of companies such as Pharmacia Biotech (Piscataway NJ), Promega (Madison WI), and US Biochemical Corp (Cleveland OH) supply commercial kits and protocols for these procedures. Suitable reporter molecules or labels include those radionuclides, enzymes,
20 fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles and the like. Patents teaching the use of such labels include US Patents 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149 and 4,366,241. Also, recombinant immunoglobulins may be produced as shown in US Patent No. 4,816,567 incorporated herein by reference.

25 **Purification of HCP**

Host cells transformed with a nucleotide sequence encoding HCP may be cultured under conditions suitable for the expression and recovery of the encoded protein from cell culture. The protein produced by a recombinant cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression
30 vectors containing an HCP-encoding sequence can be designed with signal sequences which direct secretion of HCP through a prokaryotic or eukaryotic cell membrane. Other recombinant constructions may join the sequence encoding HCP to nucleotide sequence encoding a

polypeptide domain which will facilitate purification of soluble proteins (Kroll DJ et al (1993) DNA Cell Biol 12:441-53; cf discussion of vectors infra containing fusion proteins).

HCP may also be expressed as a recombinant protein with one or more additional polypeptide domains added to facilitate protein purification. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp, Seattle WA). The inclusion of a cleavable linker sequences such as Factor XA or enterokinase (Invitrogen, San Diego CA) between the purification domain and HCP is useful to facilitate purification. One such expression vector provides for expression of a fusion protein comprising the sequence encoding HCP and nucleic acid sequence encoding 6 histidine residues followed by thioredoxin and an enterokinase cleavage site. The histidine residues facilitate purification while the enterokinase cleavage site provides a means for purifying HCP from the fusion protein.

In addition to recombinant production, fragments of HCP may be produced by direct peptide synthesis using solid-phase techniques (cf Stewart et al (1969) Solid-Phase Peptide Synthesis, WH Freeman Co, San Francisco; Merrifield J (1963) J Am Chem Soc 85:2149-2154). In vitro protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer, Foster City CA) in accordance with the instructions provided by the manufacturer. Various fragments of HCP may be chemically synthesized separately and combined using chemical methods to produce the full length molecule.

Uses of HCP

HCPs play a role in processes that involve proteolysis of specific proteins and tissues in local microenvironments including inflammation, metastasis and peptide and proenzyme processing. Since the increased expression and differential regulation of these proteases is linked to the metastatic potential of a variety of cancers, HCPs can be used to intervene in the metastatic process. Similarly, the level and effect of the cathepsins associated with protein processing diseases such as Alzheimer's disease, Huntington's disease, mucopolidosis and arthritic inflammation may be modulated to ameliorate the progression of these diseases.

A therapeutic composition comprising of antagonists, inhibitors, anti-sense molecules or anti-HCP antibodies may have application in the prevention and treatment of individuals

susceptible to diseases characterized by excessive HCPs activity. Examples include, but are not limited to, tumors, arthritic inflammations, abnormal protein processing in brain diseases and abnormal processing of regulatory proteins. A therapeutic composition comprised of agonists which modulate the activity of HCPs may have applications in diseases characterized by

5 cathepsin deficiencies including, but not limited to, mucopolipidosis I and II.

In another embodiment of the present invention, HCPs may be used to localize and quantitate the expression of HCPs with HCPs specific antibodies and nucleic acid probes based on their sequences has specific prognostic value in a variety of diseases. The correlation between high concentrations of HCPs in primary tumor cells and further occurrence of metastasis would

10 allow the identification of specific node negative breast cancer patients where chemotherapy is indicated. This prognostic potential of HCPs can also be applied to cancers of other organs including, but not limited to, the prostate and bladder.

HCP Antibodies

HCP-specific antibodies are useful for the diagnosis and treatment of conditions and

15 diseases associated with expression of HCP. Such antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, Fab fragments and fragments produced by a Fab expression library. Neutralizing antibodies, i.e., those which inhibit dimer formation, are especially preferred for diagnostics and therapeutics.

The portion of HCP used for antibody induction does not need to have biological activity;

20 however, it must be antigenic. Peptides used to induce specific antibodies may have an amino acid sequence consisting of at least five amino acids, and preferably at least 10 amino acids. Preferably, they should mimic a portion of the amino acid sequence of the natural protein and may contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of HCP amino acids may be fused with those of another protein such as keyhole limpet

25 hemocyanin and antibody produced against the chimeric molecule. Procedures well known in the art can be used for the production of antibodies to HCP

For the production of antibodies, various hosts including goats, rabbits, rats, mice, etc may be immunized by injection with HCP or any portion, fragment or oligopeptide which retains immunogenic properties. Depending on the host species, various adjuvants may be used to

30 increase immunological response. Such adjuvants include but are not limited to Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and

dinitrophenol. BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are potentially useful human adjuvants.

Monoclonal antibodies to HCP may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include but are not limited to the hybridoma technique originally described by Koehler and Milstein (1975 Nature 256:495-497), the human B-cell hybridoma technique (Kosbor et al (1983) Immunol Today 4:72; Cote et al (1983) Proc Natl Acad Sci 80:2026-2030) and the EBV-hybridoma technique (Cole et al (1985) Monoclonal Antibodies and Cancer Therapy, Alan R Liss Inc, New York NY, pp 77-96).

10 In addition, techniques developed for the production of "chimeric antibodies", the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity can be used (Morrison et al (1984) Proc Natl Acad Sci 81:6851-6855; Neuberger et al (1984) Nature 312:604-608; Takeda et al (1985) Nature 314:452-454). Alternatively, techniques described for the production of single chain antibodies
15 (US Patent No. 4,946,778) can be adapted to produce HCP-specific single chain antibodies

Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening recombinant immunoglobulin libraries or panels of highly specific binding reagents as disclosed in Orlandi et al (1989, Proc Natl Acad Sci 86: 3833-3837), and Winter G and Milstein C (1991; Nature 349:293-299).

20 Antibody fragments which contain specific binding sites for HCP may also be generated. For example, such fragments include, but are not limited to, the F(ab')₂ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab
25 fragments with the desired specificity (Huse WD et al (1989) Science 256:1275-1281).

A variety of protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the formation of complexes between HCP and its specific antibody and the measurement of complex formation. A two-site, monoclonal-based
30 immunoassay utilizing monoclonal antibodies reactive to two noninterfering epitopes on a specific HCP protein is preferred, but a competitive binding assay may also be employed. These assays are described in Maddox DE et al (1983, J Exp Med 158:1211).

Diagnostic Assays Using HCP Specific Antibodies

Particular HCP antibodies are useful for the diagnosis of conditions or diseases characterized by expression of HCP or in assays to monitor patients being treated with HCP, its fragments, agonists, antagonists or inhibitors. Diagnostic assays for HCP include methods
5 utilizing the antibody and a label to detect HCP in human body fluids or extracts of cells or tissues. The polypeptides and antibodies of the present invention may be used with or without modification. Frequently, the polypeptides and antibodies will be labeled by joining them, either covalently or noncovalently, with a reporter molecule. A wide variety of reporter molecules are known, several of which were described above.

10 A variety of protocols for measuring HCP, using either polyclonal or monoclonal antibodies specific for the respective proteins are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and fluorescent activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on HCP is preferred, but a competitive
15 binding assay may be employed. These assays are described, among other places, in Maddox, DE et al (1983, J Exp Med 158:1211).

In order to provide a basis for diagnosis, normal or standard values for HCP expression must be established. This is accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with antibody to HCP under conditions suitable for
20 complex formation which are well known in the art. The amount of standard complex formation may be quantified by comparing various artificial membranes containing known quantities of HCP with both control and disease samples from biopsied tissues. Then, standard values obtained from normal samples may be compared with values obtained from samples from subjects potentially affected by disease. Deviation between standard and subject values
25 establishes the presence of disease state.

Drug Screening

HCP, its catalytic or immunogenic fragments or oligopeptides thereof, can be used for screening therapeutic compounds in any of a variety of drug screening techniques. The fragment employed in such a test may be free in solution, affixed to a solid support, borne on a cell surface,
30 or located intracellularly. The formation of binding complexes, between HCP and the agent being tested, may be measured.

Another technique for drug screening which may be used for high throughput screening of

compounds having suitable binding affinity to the HCP is described in detail in "Determination of Amino Acid Sequence Antigenicity" by Geysen HN, WO Application 84/03564, published on September 13, 1984, and incorporated herein by reference. In summary, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or
5 some other surface. The peptide test compounds are reacted with fragments of HCP and washed. Bound HCP is then detected by methods well known in the art. Substantially purified HCP can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

10 This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding HCP specifically compete with a test compound for binding HCP. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with HCP.

Uses of the Polynucleotide Encoding HCP

15 A polynucleotide sequence encoding HCP or any part thereof may be used for diagnostic and/or therapeutic purposes. For diagnostic purposes, the sequence encoding HCP of this invention may be used to detect and quantitate gene expression in biopsied tissues in which HCP may be expressed. The diagnostic assay is useful to distinguish between absence, presence, and excess expression of HCP and to monitor regulation of HCP levels during therapeutic
20 intervention. Included in the scope of the invention are oligonucleotide sequences, antisense RNA and DNA molecules, and PNAs.

Another aspect of the subject invention is to provide for hybridization or PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding HCP or closely related molecules. The specificity of the probe, whether it is made from a highly
25 specific region, eg, 10 unique nucleotides in the 5' regulatory region, or a less specific region, eg, especially in the 3' region, and the stringency of the hybridization or amplification (maximal, high, intermediate or low) will determine whether the probe identifies only naturally occurring HCP, alleles or related sequences.

Probes may also be used for the detection of related sequences and should preferably
30 contain at least 50% of the nucleotides from any of these HCP-encoding sequences. The hybridization probes of the subject invention may be derived from the nucleotide sequence of SEQ ID NO:2 and SEQ ID NO:4 or from genomic sequence including promoter, enhancer

elements and introns of the naturally occurring sequences encoding HCP. Hybridization probes may be labeled by a variety of reporter groups, including radionuclides such as ^{32}P or ^{35}S , or enzymatic labels such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

- 5 Other means for producing specific hybridization probes for DNAs include the cloning of nucleic acid sequences encoding HCP or HCP derivatives into vectors for the production of mRNA probes. Such vectors are known in the art and are commercially available and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerase as T7 or SP6 RNA polymerase and the appropriate radioactively labeled nucleotides.

10 **Diagnostic Use**

Polynucleotide sequences encoding HCP may be used for the diagnosis of conditions or diseases with which the expression of HCP is associated. For example, polynucleotide sequences encoding HCP may be used in hybridization or PCR assays of fluids or tissues from biopsies to detect HCP expression. The form of such qualitative or quantitative methods may include

- 15 Southern or northern analysis, dot blot or other membrane-based technologies; PCR technologies; dip stick, pin, chip and ELISA technologies. All of these techniques are well known in the art and are the basis of many commercially available diagnostic kits.

The HCP-encoding nucleotide sequences disclosed herein provide the basis for assays that detect activation or induction associated with inflammation or disease. The nucleotide
20 sequence may be labeled by methods known in the art and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After an incubation period, the sample is washed with a compatible fluid which optionally contains a dye (or other label requiring a developer) if the nucleotide has been labeled with an enzyme. After the compatible fluid is rinsed off, the dye is quantitated and compared with a standard. If the
25 amount of dye in the biopsied or extracted sample is significantly elevated over that of a comparable control sample, the nucleotide sequence has hybridized with nucleotide sequences in the sample, and the presence of elevated levels of nucleotide sequences encoding HCP in the sample indicates the presence of the associated inflammation and/or disease.

Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment
30 regime in animal studies, in clinical trials, or in monitoring the treatment of an individual patient. In order to provide a basis for the diagnosis of disease, a normal or standard profile for HCP expression must be established. This is accomplished by combining body fluids or cell extracts

taken from normal subjects, either animal or human, with HCP, or a portion thereof, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained for normal subjects with a dilution series of HCP run in the same experiment where a known amount of substantially purified HCP is used. Standard values
5 obtained from normal samples may be compared with values obtained from samples from patients affected by HCP-associated diseases. Deviation between standard and subject values establishes the presence of disease.

Once disease is established, a therapeutic agent is administered and a treatment profile is generated. Such assays may be repeated on a regular basis to evaluate whether the values in the
10 profile progress toward or return to the normal or standard pattern. Successive treatment profiles may be used to show the efficacy of treatment over a period of several days or several months.

PCR, may be used as described in US Patent Nos. 4,683,195 and 4,965,188 provides additional uses for oligonucleotides based upon the sequences encoding HCP. Such oligomers are generally chemically synthesized, but they may be generated enzymatically or produced from
15 a recombinant source. Oligomers generally comprise two nucleotide sequences, one with sense orientation (5'→3') and one with antisense (3'←5'), employed under optimized conditions for identification of a specific gene or condition. The same two oligomers, nested sets of oligomers, or even a degenerate pool of oligomers may be employed under less stringent conditions for detection and/or quantitation of closely related DNA or RNA sequences.

20 Additionally, methods which may be used to quantitate the expression of a particular molecule include radiolabeling (Melby PC et al 1993 J Immunol Methods 159:235-44) or biotinylating (Duplaa C et al 1993 Anal Biochem 229-36) nucleotides, coamplification of a control nucleic acid, and standard curves onto which the experimental results are interpolated. Quantitation of multiple samples may be speeded up by running the assay in an ELISA format
25 where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation. A definitive diagnosis of this type may allow health professionals to begin aggressive treatment and prevent further worsening of the condition. Similarly, further assays can be used to monitor the progress of a patient during treatment. Furthermore, the nucleotide sequences disclosed herein may be used in molecular biology
30 techniques that have not yet been developed, provided the new techniques rely on properties of nucleotide sequences that are currently known such as the triplet genetic code, specific base pair interactions, and the like.

Therapeutic Use

Based upon the homology between HCP-1 and GI 555662 and HCP-2 and GI 181180 and their expression profiles, the polynucleotide encoding HCPs disclosed herein may be useful in the treatment of diseases including cancers and inflammation and in diseases with components
5 involving protein processing.

Expression vectors derived from retroviruses, adenovirus, herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue or cell population. Methods which are well known to those skilled in the art can be used to construct recombinant vectors which will express antisense of the sequence encoding
10 HCP. See, for example, the techniques described in Sambrook et al (supra) and Ausubel et al (supra).

The polynucleotides comprising full length cDNA sequence and/or its regulatory elements enable researchers to use the sequence encoding HCP as an investigative tool in sense (Youssoufian H and HF Lodish 1993 Mol Cell Biol 13:98-104) or antisense (Eguchi et al (1991)
15 Annu Rev Biochem 60:631-652) regulation of gene function. Such technology is now well known in the art, and sense or antisense oligomers, or larger fragments, can be designed from various locations along the coding or control regions.

Genes encoding HCP can be turned off by transfecting a cell or tissue with expression vectors which express high levels of a desired HCP fragment. Such constructs can flood cells
20 with untranslatable sense or antisense sequences. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until all copies are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector (Mettler I, personal communication) and even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing antisense molecules, DNA, RNA or PNA, to the control regions of the sequence encoding HCP, ie, the promoters, enhancers, and introns. Oligonucleotides derived from the transcription initiation site, eg, between -10 and +10 regions of the leader sequence, are preferred. The antisense molecules may also be designed to block translation of mRNA by preventing the
30 transcript from binding to ribosomes. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing compromises the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules.

Recent therapeutic advances using triplex DNA were reviewed by Gee JE et al (In: Huber BE and BI Carr (1994) Molecular and Immunologic Approaches, Futura Publishing Co, Mt Kisco NY).

Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the
5 ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage.

Within the scope of the invention are engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of the sequence encoding HCP.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites which include the following
10 sequences, GUA, GUU and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target gene containing the cleavage site may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

15 Antisense molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of RNA molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding HCP. Such DNA sequences may be incorporated into a wide variety of
20 vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, antisense cDNA constructs that synthesize antisense RNA constitutively or inducibly can be introduced into cell lines, cells or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3'
25 ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine and wybutosine as well as acetyl-, methyl-, thio- and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by
30 endogenous endonucleases.

Methods for introducing vectors into cells or tissues include those methods discussed infra and which are equally suitable for in vivo, in vitro and ex vivo therapy. For ex vivo

therapy, vectors are introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient is presented in US Patent Nos. 5,399,493 and 5,437,994, disclosed herein by reference. Delivery by transfection and by liposome are quite well known in the art.

- 5 Furthermore, the nucleotide sequences encoding HCP disclosed herein may be used in molecular biology techniques that have not yet been developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including but not limited to such properties as the triplet genetic code and specific base pair interactions.

Detection and Mapping of Related Polynucleotide Sequences

- 10 The nucleic acid sequences encoding HCP can also be used to generate hybridization probes for mapping the naturally occurring genomic sequence. The sequence may be mapped to a particular chromosome or to a specific region of the chromosome using well known techniques. These include in situ hybridization to chromosomal spreads, flow-sorted chromosomal preparations, or artificial chromosome constructions such as yeast artificial chromosomes,
15 bacterial artificial chromosomes, bacterial P1 constructions or single chromosome cDNA libraries as reviewed in Price CM (1993; Blood Rev 7:127-34) and Trask BJ (1991; Trends Genet 7:149-54).

- The technique of fluorescent in situ hybridization of chromosome spreads has been described, among other places, in Verma et al (1988) Human Chromosomes: A Manual of Basic
20 Techniques, Pergamon Press, New York NY. Fluorescent in situ hybridization of chromosomal preparations and other physical chromosome mapping techniques may be correlated with additional genetic map data. Examples of genetic map data can be found in the 1994 Genome Issue of Science (265:1981f). Correlation between the location of a the sequence encoding HCP on a physical chromosomal map and a specific disease (or predisposition to a specific disease)
25 may help delimit the region of DNA associated with that genetic disease. The nucleotide sequences of the subject invention may be used to detect differences in gene sequences between normal, carrier or affected individuals.

- In situ hybridization of chromosomal preparations and physical mapping techniques such as linkage analysis using established chromosomal markers are invaluable in extending genetic
30 maps. A recent example of an STS based map of the human genome was recently published by the Whitehead-MIT Center for Genomic Research (Hudson TJ et al (1995) Science 270:1945-1954). Often the placement of a gene on the chromosome of another mammalian

species such as mouse (Whitehead Institute/MIT Center for Genome Research, Genetic Map of the Mouse, Database Release 10, April 28, 1995) may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms, or parts thereof, by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once a disease or syndrome, such as ataxia telangiectasia (AT), has been crudely localized by genetic linkage to a particular genomic region, for example, AT to 11q22-23 (Gatti et al (1988) Nature 336:577-580), any sequences mapping to that area may represent associated or regulatory genes for further investigation. The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc. among normal, carrier or affected individuals.

Pharmaceutical Compositions

The present invention relates to pharmaceutical compositions which may comprise nucleotides, proteins, antibodies, agonists, antagonists, or inhibitors, alone or in combination with at least one other agent, such as stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water. Any of these molecules can be administered to a patient alone, or in combination with other agents, drugs or hormones, in pharmaceutical compositions where it is mixed with excipient(s) or pharmaceutically acceptable carriers. In one embodiment of the present invention, the pharmaceutically acceptable carrier is pharmaceutically inert.

Administration of Pharmaceutical Compositions

Administration of pharmaceutical compositions is accomplished orally or parenterally. Methods of parenteral delivery include topical, intra-arterial (directly to the tumor), intramuscular, subcutaneous, intramedullary, intrathecal, intraventricular, intravenous, intraperitoneal, or intranasal administration. In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of "Remington's Pharmaceutical Sciences" (Maack Publishing Co, Easton PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral

administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combination of active
5 compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; and gums
10 including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

Dragee cores are provided with suitable coatings such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene
15 glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, ie, dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating such as glycerol or sorbitol.
20 Push-fit capsules can contain active ingredients mixed with a filler or binders such as lactose or starches, lubricants such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations for parenteral administration include aqueous solutions of
25 active compounds. For injection, the pharmaceutical compositions of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active
30 compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Optionally, the suspension may also contain suitable

stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

5 **Manufacture and Storage**

The pharmaceutical compositions of the present invention may be manufactured in a manner that known in the art, eg, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many
10 acids, including but not limited to hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder in 1mM-50 mM histidine, 0.1%-2% sucrose, 2%-7% mannitol at a pH range of 4.5 to 5.5 that is combined with buffer prior to use.

15 After pharmaceutical compositions comprising a compound of the invention formulated in a acceptable carrier have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of HCP, such labeling would include amount, frequency and method of administration.

Therapeutically Effective Dose

20 Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in
25 cell culture assays, eg, of neoplastic cells, or in animal models, usually mice, rabbits, dogs, or pigs. The animal model is also used to achieve a desirable concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of protein or its antibodies,
30 antagonists, or inhibitors which ameliorate the symptoms or condition. Therapeutic efficacy and toxicity of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, eg, ED50 (the dose therapeutically effective in 50% of the

population) and LD50 (the dose lethal to 50% of the population). The dose ratio between therapeutic and toxic effects is the therapeutic index, and it can be expressed as the ratio, LD50/ED50. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of
5 dosage for human use. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

The exact dosage is chosen by the individual physician in view of the patient to be
10 treated. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Additional factors which may be taken into account include the severity of the disease state, eg, tumor size and location; age, weight and gender of the patient; diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long acting pharmaceutical compositions might be administered
15 every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from 0.1 to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature. See US Patent Nos. 4,657,760; 5,206,344; or
20 5,225,212. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

The examples below are provided to illustrate the subject invention and are not included for the purpose of limiting the invention.

25

INDUSTRIAL APPLICABILITY

I cDNA Library Construction

HCP-1

The cDNA library FIBRAGT02 was constructed from fibroblast cell lines derived from skin explants taken from the triceps area of an ataxia telangiectasia (AT) patient. The explants
30 were cultured, tested for contamination and stored as described in Weeks et al. (1991) Rad Res 128:90-99.

Samples from each line were cultured for 14 passages and treated in late log growth phase

(5 days post 1:3 split) as a monolayer in 150mm petri plates. Because AT cells are highly sensitive to radiation, equivalent dosages were determined experimentally for each cell line: D10=126+5cGy for AT fibroblasts; D10=563+31cGy for normal fibroblasts.

For normal fibroblasts, about 8.25×10^7 cells were sham-irradiated and then incubated for 5 min (22 plates) or 30 min (23 plates) at 37°C. These two sets of untreated or control cells were combined and harvested. Five minute post-gamma irradiation treatments consisted of exposing 43 plates containing about 8.49×10^7 normal cells to 50cGy from a Cs137 source and incubating them at 37°C for five minutes before harvesting. Thirty minute post-gamma irradiation treatments consisted of exposing 43 plates containing about 8.36×10^7 normal cells to 50cGy from the Cs137 source and incubating them at 37°C for thirty minutes before harvesting.

For AT fibroblasts, about 9.45×10^7 cells, were sham-irradiated and then incubated for 5 min (22 plates) or 30 min (23 plates) at 37°C. These two sets of plates (hereafter known as untreated or control AT cells) were combined and harvested. Five minute post-gamma irradiation treatments consisted of exposing 43 plates containing about 1.07×10^8 AT cells to 50cGy from a Cs137 source and incubating them at 37°C for five minutes before harvesting. Thirty minute post-gamma irradiation treatments consisted of exposing 43 plates containing about 9.71×10^7 AT cells to 50cGy from the Cs137 source and incubating them at 37°C for thirty minutes before harvesting.

The RNA was prepared by the hot phenol method and enriched for polyadenylated (poly-A) RNA by oligo d(T)-cellulose chromatography (Godbout et al. (1988) Mol Cell Biol 8:1169-1178). The yields varied from 3.0 to 5.0 percent of the total RNA. The poly-A RNA was visualized on an agarose gel, quantified, and sent to Stratagene (La Jolla CA) where the cDNA libraries were constructed in the Uni-ZAP™ vector system.

HCP-2

The cDNA library LUNGN02 was constructed from the normal lung tissue of a 47 year old male Caucasian (HEV082). The tissue was obtained from Keystone Skin Bank, International Institute for the Advancement of Medicine (Exton, PA). The frozen tissue was ground in a mortar and pestle and lysed immediately in a buffer containing guanidinium isothiocyanate. The lysate was extracted twice with phenol chloroform at pH 8.0 and centrifuged over a CsCl cushion using an Beckman SW28 rotor in a Beckman L8-70M Ultracentrifuge (Beckman Instruments). The RNA was precipitated using 0.3 M sodium acetate and 2.5 volumes of ethanol, resuspended in water and DNase treated for 15 min at 37°C. The poly A+ RNA was

isolated using the Qiagen Oligotex kit (QIAGEN Inc, Chatsworth CA).

For both FIBRAGT02 and LUNGNOT02, first strand cDNA synthesis was accomplished using an oligo d(T) primer/linker which also contained an XhoI restriction site. Second strand synthesis was performed using a combination of DNA polymerase I, *E. coli* ligase and RNase H, followed by the addition of an EcoRI adaptor to the blunt ended cDNA. The EcoRI adapted, double-stranded cDNA was then digested with XhoI restriction enzyme and fractionated on Sephacryl S400 to obtain sequences which exceeded 1000 bp in size. The size selected cDNAs were inserted into the UniZap® vector system (Stratagene, La Jolla CA); and the vector, which contains the pBluescript™ phagemid (Stratagene), was transformed into cells of *E. coli*, strain XL1-BlueMRF™ (Stratagene).

The phagemid forms of individual cDNA clones were obtained by the in vivo excision process. Enzymes from both pBluescript and a cotransformed f1 helper phage nicked the DNA, initiated new DNA synthesis, and created the smaller, single-stranded circular phagemid DNA molecules which contained the cDNA insert. The phagemid DNA was released, purified, and used to reinfect fresh host cells (SOLR, Stratagene). Presence of the phagemid which carries the gene for β -lactamase allowed transformed bacteria to grow on medium containing ampicillin.

II Isolation and Sequencing of cDNA Clones

HCP-1

Phagemid DNA for the FIBRAGT02 library was purified using the MAGIC MINIPREPS™ DNA purification system (catalogue #A7100, Promega Corp., Madison WI).

This small-scale process provides a simple and reliable method for lysing the bacterial cells and rapidly isolating purified phagemid DNA using a proprietary DNA-binding resin. The DNA was eluted from the purification resin already prepared for DNA sequencing and other analytical manipulations.

Alternatively, phagemid DNA may be purified using the QIAWELL-8 Plasmid, QIAWELL PLUS and QIAWELL ULTRA DNA purification systems (QIAGEN, Chatsworth, CA). The DNA was eluted from the purification resin and prepared for DNA sequencing and other analytical manipulations.

Plasmid DNA for the LUNGNOT02 library was purified using the Miniprep Kit (Catalogue # 77468, Advanced Genetic Technologies Corporation, Gaithersburg MD), a 96-well block kit with reagents for 960 purifications. The recommended protocol included with the kit was employed except for the following changes. Each of the 96 wells was filled with only 1 ml

of sterile Terrific Broth (Catalog # 22711, LIFE TECHNOLOGIES™) with carbenicillin at 25 mg/L and glycerol at 0.4%. After the wells were inoculated, the bacteria were cultured for 24 hours and lysed with 60 μ l of lysis buffer. A centrifugation step (Beckman GS-6R @2900 rpm for 5 min; Beckman Instruments) was performed before the contents of the block were added to the primary filter plate. The optional step of adding isopropanol to TRIS buffer was not routinely performed. After the last step in the protocol, samples were transferred to a Beckman 96-well block for storage.

The cDNAs were sequenced by the method of Sanger F and AR Coulson (1975; J Mol Biol 94:441f), using a Hamilton Micro Lab 2200 (Hamilton, Reno NV) in combination with four Peltier Thermal Cyclers (PTC200 from MJ Research, Watertown MA) and Applied Biosystems 377 or 373 DNA Sequencing Systems (Perkin Elmer) and reading frame was determined.

III Homology Searching of cDNA Clones and Their Deduced Proteins

Each cDNA was compared to sequences in GenBank using a search algorithm developed by Applied Biosystems and incorporated into the INHERIT- 670 Sequence Analysis System. In this algorithm, Pattern Specification Language (TRW Inc, Los Angeles CA) was used to determine regions of homology. The three parameters that determine how the sequence comparisons run were window size, window offset, and error tolerance. Using a combination of these three parameters, the DNA database was searched for sequences containing regions of homology to the query sequence, and the appropriate sequences were scored with an initial value. Subsequently, these homologous regions were examined using dot matrix homology plots to distinguish regions of homology from chance matches. Smith-Waterman alignments were used to display the results of the homology search.

Peptide and protein sequence homologies were ascertained using the INHERIT™ 670 Sequence Analysis System in a way similar to that used in DNA sequence homologies. Pattern Specification Language and parameter windows were used to search protein databases for sequences containing regions of homology which were scored with an initial value. Dot-matrix homology plots were examined to distinguish regions of significant homology from chance matches.

BLAST, which stands for Basic Local Alignment Search Tool (Altschul SF (1993) J Mol Evol 36:290-300; Altschul, SF et al (1990) J Mol Biol 215:403-10), was used to search for local sequence alignments. BLAST produces alignments of both nucleotide and amino acid sequences to determine sequence similarity. Because of the local nature of the alignments, BLAST is

especially useful in determining exact matches or in identifying homologs. BLAST is useful for matches which do not contain gaps. The fundamental unit of BLAST algorithm output is the High-scoring Segment Pair (HSP).

An HSP consists of two sequence fragments of arbitrary but equal lengths whose alignment is locally maximal and for which the alignment score meets or exceeds a threshold or cutoff score set by the user. The BLAST approach is to look for HSPs between a query sequence and a database sequence, to evaluate the statistical significance of any matches found, and to report only those matches which satisfy the user-selected threshold of significance. The parameter E establishes the statistically significant threshold for reporting database sequence matches. E is interpreted as the upper bound of the expected frequency of chance occurrence of an HSP (or set of HSPs) within the context of the entire database search. Any database sequence whose match satisfies E is reported in the program output.

IV Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound (Sambrook et al. supra).

Analogous computer techniques using BLAST (Altschul SF 1993 and 1990, supra) are used to search for identical or related molecules in nucleotide databases such as GenBank or the LIFESEQ™ database (Incyte, Palo Alto CA). This analysis is much faster than multiple, membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or homologous.

The basis of the search is the product score which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

and it takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1-2% error; and at 70, the match will be exact. Homologous molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

The results of the search are reported as a list of libraries in which the full length sequence, or parts thereof, is represented, the abundance of the sequence, and the percent abundance. Abundance directly reflects the number of times a particular transcript is present in a

cDNA library, and percent abundance is abundance divided by the number of sequences examined in the library.

V Extension of the Sequences Encoding HCP

The nucleic acid sequence of SEQ ID NO:2 and SEQ ID NO:4 is used to design oligo-
 5 nucleotide primers for extending a partial nucleotide sequence to full length or for obtaining
 5' sequence from genomic libraries. One primer is synthesized to initiate extension in the
 antisense direction (XLR) and the other is synthesized to extend sequence in the sense direction
 (XLF). Primers allow the extension of the known sequence "outward" generating amplicons
 containing new, unknown nucleotide sequence for the region of interest (US Patent Application
 10 08/487,112, filed June 7, 1995, specifically incorporated by reference). The initial primers are
 designed from the cDNA using OLIGO® 4.06 Primer Analysis Software (National Biosciences),
 or another appropriate program, to be 22-30 nucleotides in length, to have a GC content of 50%
 or more, and to anneal to the target sequence at temperatures about 68°-72° C. Any stretch of
 nucleotides which would result in hairpin structures and primer-primer dimerizations is avoided.
 15 The original, selected cDNA libraries, or a human genomic library are used to extend the
 sequence; the latter is most useful to obtain 5' upstream regions. If more extension is necessary
 or desired, additional sets of primers are designed to further extend the known region.

By following the instructions for the XL-PCR kit (Perkin Elmer) and thoroughly mixing
 the enzyme and reaction mix, high fidelity amplification is obtained. Beginning with 40 pmol of
 20 each primer and the recommended concentrations of all other components of the kit, PCR is
 performed using the Peltier Thermal Cycler (PTC200; MJ Research, Watertown MA) and the
 following parameters:

	Step 1	94° C for 1 min (initial denaturation)
	Step 2	65° C for 1 min
25	Step 3	68° C for 6 min
	Step 4	94° C for 15 sec
	Step 5	65° C for 1 min
	Step 6	68° C for 7 min
	Step 7	Repeat step 4-6 for 15 additional cycles
30	Step 8	94° C for 15 sec
	Step 9	65° C for 1 min
	Step 10	68° C for 7:15 min
	Step 11	Repeat step 8-10 for 12 cycles
	Step 12	72° C for 8 min
35	Step 13	4° C (and holding)

A 5-10 µl aliquot of the reaction mixture is analyzed by electrophoresis on a low

concentration (about 0.6-0.8%) agarose mini-gel to determine which reactions were successful in extending the sequence. Bands thought to contain the largest products were selected and cut out of the gel. Further purification involves using a commercial gel extraction method such as QIAQuick™ (QIAGEN Inc). After recovery of the DNA, Klenow enzyme was used to trim
 5 single-stranded, nucleotide overhangs creating blunt ends which facilitate religation and cloning.

After ethanol precipitation, the products are redissolved in 13 μ l of ligation buffer, 1 μ l T4-DNA ligase (15 units) and 1 μ l T4 polynucleotide kinase are added, and the mixture is incubated at room temperature for 2-3 hours or overnight at 16° C. Competent *E. coli* cells (in 40 μ l of appropriate media) are transformed with 3 μ l of ligation mixture and cultured in 80 μ l of
 10 SOC medium (Sambrook J et al, supra). After incubation for one hour at 37° C, the whole transformation mixture is plated on Luria Bertani (LB)-agar (Sambrook J et al, supra) containing 2xCarb. The following day, several colonies are randomly picked from each plate and cultured in 150 μ l of liquid LB/2xCarb medium placed in an individual well of an appropriate, commercially-available, sterile 96-well microtiter plate. The following day, 5 μ l of each
 15 overnight culture is transferred into a non-sterile 96-well plate and after dilution 1:10 with water, 5 μ l of each sample is transferred into a PCR array.

For PCR amplification, 18 μ l of concentrated PCR reaction mix (3.3x) containing 4 units of rTth DNA polymerase, a vector primer and one or both of the gene specific primers used for the extension reaction are added to each well. Amplification is performed using the following
 20 conditions:

Step 1	94° C for 60 sec
Step 2	94° C for 20 sec
Step 3	55° C for 30 sec
Step 4	72° C for 90 sec
25 Step 5	Repeat steps 2-4 for an additional 29 cycles
Step 6	72° C for 180 sec
Step 7	4° C (and holding)

Aliquots of the PCR reactions are run on agarose gels together with molecular weight markers. The sizes of the PCR products are compared to the original partial cDNAs, and
 30 appropriate clones are selected, ligated into plasmid and sequenced.

VI Labeling and Use of Hybridization Probes

Hybridization probes derived from SEQ ID NO:2 are employed to screen cDNAs, genomic DNAs or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base-pairs, is specifically described, essentially the same procedure is used with larger cDNA

fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 (National Biosciences), labeled by combining 50 pmol of each oligomer and 250 mCi of [γ - 32 P] adenosine triphosphate (Amersham, Chicago IL) and T4 polynucleotide kinase (DuPont NEN®, Boston MA). The labeled oligonucleotides are substantially purified with Sephadex G-25 super fine resin column (Pharmacia). A portion containing 10^7 counts per minute of each of the sense and antisense oligonucleotides is used in a typical membrane based hybridization analysis of human genomic DNA digested with one of the following endonucleases (Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II; DuPont NEN®).

The DNA from each digest is fractionated on a 0.7 percent agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under increasingly stringent conditions up to 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. After XOMAT AR™ film (Kodak, Rochester NY) is exposed to the blots in a Phosphorimager cassette (Molecular Dynamics, Sunnyvale CA) for several hours, hybridization patterns are compared visually.

VII Antisense Molecules

The sequence encoding HCP, or any part thereof, is used to inhibit in vivo or in vitro expression of naturally occurring sequence. Although use of antisense oligonucleotides, comprising about 20 base-pairs, is specifically described, essentially the same procedure is used with larger cDNA fragments. An oligonucleotide complementary to the coding sequence of HCP as shown in Figures 1A, 1B, 1C, 1D, 1E, 2A, 2B, 3 and 4 is used to inhibit expression of naturally occurring sequence. The complementary oligonucleotide is designed from the most unique 5' sequence as shown in Figures 1, 2, 3, and 4 and used either to inhibit transcription by preventing promoter binding to the upstream nontranslated sequence or translation of an HCP-encoding transcript by preventing the ribosome from binding. Using an appropriate portion of the leader and 5' sequence of SEQ ID NO:2 or SEQ ID NO:4 an effective antisense oligonucleotide includes any 15-20 nucleotides spanning the region which translates into the signal or early coding sequence of the polypeptide as shown in Figures 1A, 1B, 1C, 1D, 1E, 2A, 2B, 3, and 4.

VIII Expression of HCP

Expression of the HCP is accomplished by subcloning the cDNAs into appropriate vectors and transfecting the vectors into host cells. In this case, the cloning vector, pSport,

previously used for the generation of the cDNA library is used to express HCP in *E. coli*. Upstream of the cloning site, this vector contains a promoter for β -galactosidase, followed by sequence containing the amino-terminal Met and the subsequent 7 residues of β -galactosidase. Immediately following these eight residues is a bacteriophage promoter useful for transcription
5 and a linker containing a number of unique restriction sites.

Induction of an isolated, transfected bacterial strain with IPTG using standard methods produces a fusion protein which consists of the first seven residues of β -galactosidase, about 5 to 15 residues of linker, and the full length HCP. The signal sequence directs the secretion of HCP into the bacterial growth media which can be used directly in the following assay for activity.

10 IX Assay for HCP Activity

The characterization of protease activity and specificity is based on the rate of cleavage of specific peptide substrates and a determination of an inhibitor profile. Rates of cleavage for cathepsin L are assessed by incubation of the protease with substrates such as Z-Phe-Arg-AMC or Bz-Val-Lys-Lys-Arg-AFC and measuring the rate of release of the fluorescent or chromogenic
15 leaving groups. Further specificity of the protease can be examined by titrating specific inhibitors into the cleavage assays and examining the change in the rate of proteolysis. Inhibitors for cathepsin L include trans-epoxysuccinyl-L-leucylamido-(3-methyl)butane, trans-epoxysuccinyl-L-leucylamido-(4-guanidino)butane, chymostatin, and PLCPI.

Rates of cleavage for cathepsin D are assessed by incubation of the protease with
20 substrates such as D-Phe-Ser-(Bzl)-Phe-Phe-Ala-Ala-p-aminobenzoate and measuring the rate of release of the leaving group. Specificity of cathepsin D is confirmed by the effect of specific inhibitors including pepstatin A.

X Production of HCP Specific Antibodies

HCP substantially purified using PAGE electrophoresis (Sambrook, supra) are used to
25 immunize rabbits and to produce antibodies using standard protocols. The amino acid sequence translated from HCP is analyzed using DNASTar software (DNASTar Inc) to determine regions of high immunogenicity and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Analysis to select appropriate epitopes, such as those near the C-terminus or in hydrophilic regions (shown in Figures 1A, 1B, 1C, 1D,
30 1E, 4 and 7) is described by Ausubel FM et al (supra).

Typically, the oligopeptides are 15 residues in length, synthesized using an Applied Biosystems Peptide Synthesizer Model 431A using fmoc-chemistry, and coupled to keyhole

limpet hemocyanin (KLH, Sigma) by reaction with M-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS; Ausubel FM et al, supra). Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. The resulting antisera are tested for antipeptide activity, for example, by binding the peptide to plastic, blocking with 1% BSA, reacting with rabbit
5 antisera, washing, and reacting with radioiodinated, goat anti-rabbit IgG.

XI Purification of Naturally Occurring HCP Using Specific Antibodies

Naturally occurring or recombinant HCP are substantially purified by immunoaffinity chromatography using antibodies specific for HCP. An immunoaffinity column is constructed by covalently coupling HCP antibody to an activated chromatographic resin such as CnBr-activated
10 Sepharose (Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Membrane fractions from cells expressing HCP are prepared by methods well known in the art. Alternatively, a recombinant HCP fragment containing an appropriate signal sequence may be secreted in useful quantity into the medium in which transfected cells are grown.
15 An HCP-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of HCP (eg, high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/HCP binding (eg, a buffer of pH 2-3 or a high concentration of a chaotrope such as urea or thiocyanate ion), and HCP is collected.

20 All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited
25 to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION

- (i) APPLICANT: INCYTE PHARMACEUTICALS, INC.
- (ii) TITLE OF THE INVENTION: TWO NOVEL HUMAN CATHESPIN PROTEINS
- (iii) NUMBER OF SEQUENCES: 4
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Incyte Pharmaceuticals, Inc.
 - (B) STREET: 3174 Porter Drive
 - (C) CITY: Palo Alto
 - (D) STATE: CA
 - (E) COUNTRY: U.S.
 - (F) ZIP: 94304
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Diskette
 - (B) COMPUTER: IBM Compatible
 - (C) OPERATING SYSTEM: DOS
 - (D) SOFTWARE: FastSEQ Version 1.5
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: To Be Assigned
 - (B) FILING DATE: Filed Herewith
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 08/723,938
 - (B) FILING DATE: 26-SEP-1996
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Billings, Lucy J.
 - (B) REGISTRATION NUMBER: 36,749
 - (C) REFERENCE/DOCKET NUMBER: PF-0125 PCT
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 650-855-0555
 - (B) TELEFAX: 650-845-4166

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 392 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- (vii) IMMEDIATE SOURCE:
 - (A) LIBRARY:
 - (B) CLONE: Consensus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Met Val Cys Arg Leu Pro Val Ser Lys Lys Thr Leu Leu Cys Ser Phe
 1 5 10 15

```

Gln Val Leu Asp Glu Leu Gly Arg His Val Leu Leu Arg Lys Asp Cys
      20      25      30
Gly Pro Val Asp Thr Lys Val Pro Gly Ala Val Glu Pro Lys Ser Ala
      35      40      45
Phe Thr Gln Gly Ser Ala Met Ile Ser Ser Leu Ser Gln Asn His Pro
      50      55      60
Asp Asn Arg Asn Glu Thr Phe Ser Ser Xaa Ile Ser Leu Leu Asn Glu
      65      70      75      80
Asp Pro Leu Ser Gln Asp Leu Pro Val Lys Met Ala Ser Ile Phe Lys
      85      90      95
Asn Phe Val Ile Thr Tyr Asn Arg Thr Tyr Glu Ser Lys Glu Glu Ala
      100      105      110
Arg Trp Arg Leu Ser Val Phe Val Asn Asn Met Val Arg Ala Gln Lys
      115      120      125
Ile Gln Ala Leu Asp Arg Gly Thr Ala Gln Tyr Gly Val Thr Lys Phe
      130      135      140
Ser Asp Leu Thr Glu Glu Phe Arg Thr Ile Tyr Leu Asn Thr Leu
      145      150      155      160
Leu Arg Lys Glu Pro Gly Asn Lys Met Lys Gln Ala Lys Ser Val Gly
      165      170      175
Asp Leu Ala Pro Pro Glu Trp Asp Trp Arg Ser Lys Gly Ala Val Thr
      180      185      190
Lys Val Lys Asp Gln Gly Met Cys Gly Ser Cys Trp Ala Phe Ser Val
      195      200      205
Thr Gly Asn Val Glu Gly Gln Trp Phe Leu Asn Gln Gly Thr Leu Leu
      210      215      220
Ser Leu Ser Glu Gln Glu Leu Leu Asp Cys Asp Lys Met Asp Lys Ala
      225      230      235      240
Cys Met Gly Gly Leu Pro Ser Asn Ala Tyr Ser Ala Ile Lys Asn Leu
      245      250      255
Gly Gly Leu Glu Thr Glu Asp Asp Tyr Ser Tyr Gln Gly His Met Gln
      260      265      270
Ser Cys Asn Phe Ser Ala Glu Lys Ala Lys Val Tyr Ile Asn Asp Ser
      275      280      285
Val Glu Leu Ser Gln Asn Glu Gln Lys Leu Ala Ala Trp Leu Ala Glu
      290      295      300
Arg Gly Pro Ile Ser Val Ala Ile Asn Ala Phe Gly Met Gln Phe Tyr
      305      310      315      320
Arg His Gly Ile Ser Arg Pro Leu Arg Pro Leu Cys Ser Pro Trp Leu
      325      330      335
Ile Asp His Ala Val Leu Leu Val Gly Tyr Gly Asn Arg Ser Asp Val
      340      345      350
Pro Phe Trp Ala Ile Lys Asn Ser Trp Gly Thr Asp Trp Gly Glu Lys
      355      360      365
Gly Tyr Tyr Tyr Leu His Arg Gly Ser Gly Ala Cys Gly Val Asn Thr
      370      375      380
Met Ala Ser Ser Ala Val Val Asp
      385      390

```

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1642 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY:

(B) CLONE: Consensus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GGCCACCCTG	GAGGAGCCAC	CCTGGAACGA	CCCCATGGTG	TGCCGGCTCC	CCGTGTCCAA	60
GAAAACCCTG	CTCTGCAGCT	TCCAAGTCCT	GGATGAGCTC	GGAAGACACG	TGCTGCTGCG	120
GAAGGACTGT	GGCCCAAGTG	ACACCAAGGT	TCCAGGTGCT	GTGGAGCCCA	AGTCAGCCTT	180
TACTCAGGGC	TCAGCCATGA	TTTCTTCTCT	GTCCCAAAAC	CATCCAGACA	ACAGAAACGA	240
GACTTTCAGC	TCARTCATTT	CCCTGTTGAA	TGAGGATCCC	CTGTCCCAGG	ACTTGCTGTG	300
GAAGATGGCT	TCAATCTTCA	AGAACTTTGT	CATTACCTAT	AACCGGACAT	ATGAGTCAAA	360
GGAAGAAGCC	CGGTGGCGCC	TGTCCGTCTT	TGTCAATAAC	ATGGTGCGAG	CACAGAAGAT	420
CCAGGCCCTG	GACCGTGGCA	CAGCTCAGTA	TGGAGTCACC	AAGTTCAGTG	ATCTCACAGA	480
GGAGGAGTTC	CGCACTATCT	ACCTGAATAC	TCTCCTGAGA	AAAGAGCCTG	GCAACAAGAT	540
GAAGCAAGCC	AAGTCTGTGG	GTGACCTCGC	CCCACCTGAA	TGGGACTGGA	GGAGTAAGGG	600
GGCTGTCACA	AAAGTCAAAG	ACCAGGGCAT	GTGTGGCTCC	TGCTGGGCCT	TCTCAGTCAC	660
AGGCAATGTG	GAGGGCCAGT	GGTTTCTCAA	CCAGGGGACC	CTGCTCTCCC	TCTCTGAACA	720
GGAGCTCTTG	GACTGTGACA	AGATGGACAA	GGCCTGCATG	GGCGGCTTGC	CCTCCAATGC	780
CTACTCGGCC	ATAAAGAATT	TGGGAGGGCT	GGAGACAGAG	GATGACTACA	GCTACCAAGG	840
TCACATGCAG	TCCTGCAACT	TCTCAGCAGA	GAAGGCCAAG	GTCTACATCA	ATGACTCCGT	900
GGAGCTGAGC	CAGAACGAGC	AGAAGCTGGC	AGCCTGGCTG	GCCGAGAGAG	GCCCAATCTC	960
CGTGGCCATC	AATGCCTTTG	GCATGCAGTT	TTACCGCCAC	GGGATCTCCC	GCCCTCTCCG	1020
ACCCCTCTGC	AGCCCTTGGC	TCATTGACCA	TGCGGTGTTG	CTTGTGGGCT	ACGGCAACCG	1080
CTCTGACGTT	CCCTTTTGGG	CCATCAAGAA	CAGCTGGGGC	ACTGACTGGG	GTGAGAAGGG	1140
TTACTACTAC	TTGCATCGCG	GGTCCGGGGC	CTGTGGCGTG	AACACCATGG	CCAGCTCGGC	1200
GGTGGTGGAC	TGATGAGGGG	CCCCAGCTC	GGGGCCTGGT	GCTGATCAGA	GTGGCTGCTG	1260
CCCCAGCCTG	ACATGTGTCC	AGGCCCCCTC	CCGGGAGGTA	CAGCTGGCAG	AGGGAAAGGC	1320
ACTGGGTACC	TCAGGGTGAG	CAGAGGGCAC	TGGGCTGGGG	CACAGCCCCT	GCTTCCCTGC	1380
ACCCCATTC	CACCCTGAAG	TTCTGCACCT	GCACCTTTGT	TGAATTGTGG	TAGCTTAGGA	1440
GGATGTGCGG	GTTGAAGGGT	GGTATCTTGG	CAGTTGAAGC	TGGGGCAAGA	ACTCTGGGCT	1500
TGGGTAATGA	AGCAGGAAGA	AAATTTTCTT	GATCTTAAGC	CCACCTCTGT	TCTGCCCCCT	1560
GCTTTTCCTC	TGTTTGAATA	CTATAAATTT	TCTGGTTCCC	TTTGGATTTA	TGGATAAGTG	1620
TCCCCCTCCA	TGTTCCAGGA	AA				1642

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 395 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: LUNGNOT02
- (B) CLONE: 312099

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Met	Ser	Pro	Pro	Pro	Leu	Leu	Gln	Pro	Leu	Leu	Leu	Leu	Leu	Pro	Leu
1				5					10					15	
Leu	Asn	Val	Glu	Pro	Ser	Gly	Ala	Thr	Leu	Ile	Arg	Ile	Pro	Leu	His
			20					25					30		
Arg	Val	Gln	Pro	Gly	Arg	Arg	Ile	Leu	Asn	Leu	Leu	Arg	Gly	Trp	Arg
		35					40					45			
Glu	Pro	Ala	Glu	Leu	Pro	Lys	Leu	Gly	Ala	Pro	Ser	Pro	Gly	Asp	Lys
		50				55					60				
Pro	Ile	Phe	Val	Pro	Leu	Ser	Asn	Tyr	Arg	Asp	Val	Gln	Tyr	Phe	Gly
65					70					75					80
Glu	Ile	Gly	Leu	Gly	Thr	Pro	Pro	Gln	Asn	Phe	Thr	Val	Ala	Phe	Asp
			85						90					95	
Thr	Gly	Ser	Ser	Asn	Leu	Trp	Val	Pro	Ser	Arg	Arg	Cys	His	Phe	Phe
			100					105					110		
Ser	Val	Pro	Cys	Trp	Leu	His	His	Arg	Phe	Asp	Pro	Lys	Ala	Ser	Ser
		115					120					125			
Ser	Phe	Gln	Ala	Asn	Gly	Thr	Lys	Phe	Ala	Ile	Gln	Tyr	Gly	Thr	Gly
		130				135					140				
Arg	Val	Asp	Gly	Ile	Leu	Ser	Glu	Asp	Lys	Leu	Thr	Ile	Gly	Gly	Ile
145					150					155					160
Lys	Gly	Ala	Ser	Val	Ile	Phe	Gly	Glu	Ala	Leu	Trp	Glu	Pro	Ser	Leu
			165						170					175	
Val	Phe	Ala	Phe	Ala	His	Phe	Asp	Gly	Ile	Leu	Gly	Leu	Gly	Phe	Pro
			180					185					190		
Ile	Leu	Ser	Val	Glu	Gly	Val	Arg	Pro	Pro	Met	Asp	Val	Leu	Val	Glu
		195					200					205			
Gln	Gly	Leu	Leu	Asp	Lys	Pro	Val	Phe	Ser	Phe	Tyr	Leu	Asn	Arg	Asp
		210				215					220				
Pro	Glu	Glu	Pro	Asp	Gly	Gly	Glu	Leu	Val	Leu	Gly	Gly	Ser	Asp	Pro
225					230					235					240
Ala	His	Tyr	Ile	Pro	Pro	Leu	Thr	Phe	Val	Pro	Val	Thr	Val	Pro	Ala
			245						250					255	
Tyr	Trp	Gln	Ile	His	Met	Glu	Arg	Val	Lys	Val	Gly	Pro	Gly	Leu	Thr
		260						265					270		
Leu	Cys	Ala	Lys	Gly	Cys	Ala	Ala	Ile	Leu	Asp	Thr	Gly	Thr	Ser	Leu
		275					280					285			
Ile	Thr	Gly	Pro	Thr	Glu	Glu	Ile	Arg	Ala	Leu	His	Ala	Ala	Ile	Gly
		290				295					300				
Gly	Ile	Pro	Leu	Leu	Ala	Gly	Glu	Tyr	Ile	Ile	Leu	Cys	Ser	Glu	Ile
305					310					315					320
Pro	Lys	Leu	Pro	Ala	Val	Ser	Phe	Leu	Leu	Gly	Gly	Val	Trp	Phe	Asn
			325						330					335	
Leu	Thr	Ala	His	Asp	Tyr	Val	Ile	Gln	Thr	Thr	Arg	Asn	Asp	Val	Phe
			340						345				350		
Leu	Gly	Thr	Tyr	Val	Ala	Val	Phe	Asp	Arg	Gly	Asp	Met	Lys	Ser	Ser
		355					360					365			
Ala	Arg	Val	Gly	Leu	Ala	Arg	Ala	Arg	Thr	Arg	Gly	Ala	Asp	Leu	Gly
		370				375						380			
Trp	Gly	Glu	Thr	Ala	Gln	Ala	Gln	Phe	Pro	Gly					
385					390					395					

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1299 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vii) IMMEDIATE SOURCE:

(A) LIBRARY: LUNGNOT02

(B) CLONE: 312099

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CCACGCGTCC	GCAGCGATGT	CTCCACCACC	GCTGCTGCAA	CCCCTGCTGC	TGCTGCTGCC	60
TCTGCTGAAT	GTGGAGCCTT	CCGGGGCCAC	ACTGATCCGC	ATCCCTCTTC	ATCGAGTCCA	120
ACCTGGACGC	AGGATCCTGA	ACCTACTGAG	GGGATGGAGA	GAACCAGCAG	AGCTCCCCAA	180
GTTGGGGGCC	CCATCCCCTG	GGGACAAGCC	CATCTTCGTA	CCTCTCTCGA	ACTACAGGGA	240
TGTGCAGTAT	TTTGGGGAAA	TTGGGCTGGG	AACGCCTCCA	CAAAACTTCA	CTGTTGCCTT	300
TGACACTGGC	TCCTCCAATC	TCTGGGTCCC	GTCCAGGAGA	TGCCACTTCT	TCAGTGTGCC	360
CTGCTGGTTA	CACCACCGAT	TTGATCCCAA	AGCCTCTAGC	TCCTTCCAGG	CCAAATGGGAC	420
CAAGTTTGCC	ATTCAATATG	GAAGTGGGCG	GGTAGATGGA	ATCCTGAGCG	AGGACAAGCT	480
GACTATTGGT	GGAATCAAGG	GTGCATCAGT	GATTTTCGGG	GAGGCTCTCT	GGGAGCCCAG	540
CCTGGTCTTC	GCTTTTGCCC	ATTTTGATGG	GATATTGGGC	CTCGGTTTTC	CCATTCTGTC	600
TGTGGAAGGA	GTTCGGCCCC	CGATGGATGT	ACTGGTGGAG	CAGGGGCTAT	TGGATAAGCC	660
TGTCTTCTCC	TTTACCTCA	ACAGGGACCC	TGAAGAGCCT	GATGGAGGAG	AGCTGGTCCT	720
GGGGGGCTCG	GACCCGGCAC	ACTACATCCC	ACCCCTCACC	TTCGTGCCAG	TCACGGTCCC	780
CGCCTACTGG	CAGATCCACA	TGGAGCGTGT	GAAGGTGGGC	CCAGGGCTGA	CTCTCTGTGC	840
CAAGGGCTGT	GCTGCCATCC	TGGATACGGG	CACGTCCCTC	ATCACAGGAC	CCACTGAGGA	900
GATCCGGGCC	CTGCATGCAG	CCATTGGGGG	AATCCCCTTG	CTGGCTGGGG	AGTACATCAT	960
CCTGTGCTCG	GAAATCCCAA	AGCTCCCGC	AGTCTCCTTC	CTTCTTGGGG	GGGTCTGGTT	1020
TAACCTCACG	GCCCATGATT	ACGTCATCCA	GACTACTCGA	AATGACGTCT	TCTTGGGGAC	1080
GTATGTGGCC	GTCTTCGACC	GCGGGGACAT	GAAGAGCAGC	GCCCGGGTGG	GCCTGGCGCG	1140
CGCTCGCACT	CGCGGAGCGG	ACCTCGGATG	GGGAGAGACT	GCGCAGGCGC	AGTTCCCCGG	1200
GTGACGCCCA	AGTGAAGCGC	ATGCGCAGCG	GGTGGTCGCG	GAGGTCTCTG	TACCCAGTAA	1260
AAATCCACTA	TTTCATTGA	AAAAAAAAAA	AAAAAAAAAA			1299

CLAIMS

1. A substantially purified human cathepsin polypeptide comprising the amino acid sequence of SEQ ID NO:1, or fragments thereof.
2. An isolated polynucleotide sequence encoding the polypeptide of claim 1.
- 5 3. An isolated polynucleotide sequence consisting of the nucleic acid sequence of SEQ ID NO:2 or variants thereof.
4. An isolated polynucleotide sequence consisting of the complement of SEQ ID NO:2.
5. An isolated polynucleotide sequence which hybridizes under stringent conditions to SEQ ID NO:2.
- 10 6. A hybridization probe consisting of SEQ ID NO:2, or fragments thereof.
7. A recombinant expression vector containing the polynucleotide sequence of claim 3.
8. A recombinant host cell containing the vector of claim 7.
9. A method for producing the polypeptide consisting of SEQ ID NO:1, or fragments thereof, the method comprising the steps of:
 - 15 a) culturing the host cell of claim 8 under conditions suitable for the expression of the polypeptide; and
 - b) recovering the polypeptide from the host cell culture.
10. A pharmaceutical composition comprising a substantially purified human cathepsin protein having the amino acid sequence of SEQ ID NO:1 in conjunction with a suitable
20 pharmaceutical carrier.
11. A method of treating cancer comprising administering to an individual in need of such treatment an effective amount of the pharmaceutical composition of claim 10.
12. A purified antibody which binds specifically to the polypeptide of claim 1.
13. A purified agonist which specifically modulates the biological activity of the
25 polypeptide of claim 1.
14. A purified antagonist which specifically modulates the biological activity of the polypeptide of claim 1.
15. A substantially purified human cathepsin polypeptide comprising the amino acid sequence of SEQ ID NO:3, or fragments thereof.
- 30 16. An isolated polynucleotide sequence encoding the polypeptide of claim 15.
17. An isolated polynucleotide sequence consisting of the nucleic acid sequence of SEQ ID NO:4 or variants thereof.

18. An isolated polynucleotide sequence consisting of the complement of SEQ ID NO:4.
19. An isolated polynucleotide sequence which hybridizes under stringent conditions to SEQ ID NO:4.
20. A hybridization probe consisting of SEQ ID NO:4, or fragments thereof.
- 5 21. A recombinant expression vector containing the polynucleotide sequence of claim 17.
22. A recombinant host cell containing the vector of claim 21.
23. A method for producing the polypeptide consisting of SEQ ID NO:3, or fragments thereof, the method comprising the steps of:
 - a) culturing the host cell of claim 22 under conditions suitable for the expression
 - 10 of the polypeptide; and
 - b) recovering the polypeptide from the host cell culture.
24. A pharmaceutical composition comprising a substantially purified human cathepsin protein having the amino acid sequence of SEQ ID NO:3 in conjunction with a suitable pharmaceutical carrier.
- 15 25. A method of treating cancer comprising administering to an individual in need of such treatment an effective amount of the pharmaceutical composition of claim 24.
26. A purified antibody which binds specifically to the polypeptide of claim 15.
27. A purified agonist which specifically modulates the biological activity of the polypeptide of claim 15.
- 20 28. A purified antagonist which specifically modulates the biological activity of the polypeptide of claim 15.

1/17

5' GCC ACC CTG GAG GAG CCA CCC TGG AAC GAC CCC ATG GTG TGC CGG CTC CCC GTG
 9 18 27 36 45 54
 M V C R L P V

 TCC AAG AAA ACC CTG CTC TGC AGC TTC CAA GTC CTG GAT GAG CTC GGA AGA CAC
 63 72 81 90 99 108
 S K K T L L C S F Q V L D E L G R H

 GTG CTG CTG CGG AAG GAC TGT GGC CCA GTG GAC ACC AAG GTT CCA GGT GCT GTG
 117 126 135 144 153 162
 V L L R K D C G P V D T K V P G A V

 GAG CCC AAG TCA GCC TTT ACT CAG GGC TCA GCC ATG ATT TCT CTG TCC CAA
 171 180 189 198 207 216
 E P K S A F T Q G S A M I S S L S Q

 AAC CAT CCA GAC AAC AGA AAC GAG ACT TTC AGC TCA RTC ATT TCC CTG TTG AAT
 225 234 243 252 261 270
 N H P D N R N E T F S S X I S L L N

 GAG GAT CCC CTG TCC CAG GAC TTG CCT GTG AAG ATG GCT TCA ATC TTC AAG AAC
 279 288 297 306 315 324
 E D P L S Q D L P V K M A S I F K N

 TTT GTC ATT ACC TAT AAC CGG ACA TAT GAG TCA AAG GAA GAA GCC CGG TGG CGC
 333 342 351 360 369 378
 F V I T Y N R T Y E S K E A R W R

FIGURE 1A

2/17

387	CTG TCC GTC TTT GTC AAT AAC ATG GTG CGA GCA CAG AAG ATC CAG GCC CTG GAC	414	423	432
	L S V F V N N N M V R A Q K I Q A L D			
441	CGT GGC ACA GCT CAG TAT GGA GTC ACC AAG TTC AGT GAT CTC ACA GAG GAG GAG	468	477	486
	R G T A Q Y G V T K F S D L T E E E			
495	TTC CGC ACT ATC TAC CTG AAT ACT CTC CTG AGA AAA GAG CCT GGC AAC AAG ATG	522	531	540
	F R T I Y L Y L N T L L L R K E P G N K M			
549	AAG CAA GCC AAG TCT GTG GGT GAC CTC GGC CCA CCT GAA TGG GAC TGG AGG AGT	576	585	594
	K Q A K S V G D L A P P E W D W R S			
603	AAG GGG GCT GTC ACA AAA GTC AAA GAC GGC ATG TGT GGC TCC TGC TGG GCC	630	639	648
	K G A V T K V K D Q G M C G S C W A			
657	TTC TCA GTC ACA GGC AAT GTG GAG GGC CAG TGG TTT CTC AAC CAG GGG ACC CTG	684	693	702
	F S V T G N V E G Q W F L N Q G T L			
711	CTC TCC CTC TCT GAA CAG GAG CTC TTG GAC TGT GAC AAG ATG GAC AAG GCC TGC	738	747	756
	L S L S E Q E L L L D C D K M D K A C			

FIGURE 1B

3/17

765	774	783	792	801	810
ATG GGC GGC TTG CCC TCC AAT GCC TAC TCG GCC ATA AAG AAT TTG GGA GGG CTG					
M G G L P S N A Y S A I K N L G G L					
819	828	837	846	855	864
GAG ACA GAG GAT GAC TAC AGC TAC CAG GGT CAC ATG CAG TCC TGC AAC TTC TCA					
E T E D D Y S Y Q G H M Q S C N F S					
873	882	891	900	909	918
GCA GAG AAG GCC AAG GTC TAC ATC AAT GAC TCC GTG GAG CTG AGC CAG AAC GAG					
A E K A A K V Y I N D S V E L S Q N E					
927	936	945	954	963	972
CAG AAG CTG GCA GCC TGG CTG GCC GAG AGA GGC CCA ATC TCC GTG GCC ATC AAT					
Q K L A A W L A E R G P I S V A I N					
981	990	999	1008	1017	1026
GCC TTT GGC ATG CAG TTT TAC CGC CAC GGG ATC TCC CGC CCT CTC CGA CCC CTC					
A F G M Q F Y R H G I S R P L R P L					
1035	1044	1053	1062	1071	1080
TGC AGC CCT TGG CTC ATT GAC CAT GCG GTG TTG CTT GTG GGC TAC GGC AAC CGC					
C S P W L I D H A V L L V G Y G N R					
1089	1098	1107	1116	1125	1134
TCT GAC GTT CCC TTT TGG GCC ATC AAG AAC AGC TGG GGC ACT GAC TGG GGT GAG					
S D V P F W A I K N S W G T D W G E					

FIGURE 1C

4/17

1143 AAG GGT TAC TAC TAC TTG CAT CGC GGG TCC GGG GCC TGT GGC GTG AAC ACC ATG 1188
 K G Y Y Y L H R G S G A C G V N T M
 1197 1206 1215 1224 1233 1242
 GCC AGC TCG GCG GTG GTG GAC TGA TGA GGG GCC CCC AGC TCG GGG CCT GGT GCT
 A S S A V V D *
 1251 1260 1269 1278 1287 1296
 GAT CAG AGT GGC TGC TGC CCC AGC CTG ACA TGT GTC CAG GCC CCT CCC CGG GAG
 1305 1314 1323 1332 1341 1350
 GTA CAG CTG GCA GAG GGA AAG GCA CTG GGT ACC TCA GGG TGA GCA GAG GGC ACT
 1359 1368 1377 1386 1395 1404
 GGG CTG GGG CAC AGC CCC TGC TTC CCT GCA CCC CAT TCC CAC CCT GAA GTT CTG
 1413 1422 1431 1440 1449 1458
 CAC CTG CAC CTT TGT TGA ATT GTG GTA GCT TAG GAG GAT GTC GGG GTT GAA GGG
 1467 1476 1485 1494 1503 1512
 TGG TAT CTT GGC AGT TGA AGC TGG GGC AAG AAC TCT GGG CTT GGG TAA TGA AGC

FIGURE 1D

5/17

```
1521      1530      1539      1548      1557      1566
AGG AAG AAA ATT TTC TTG ATC TTA AGC CCA CCT CTG TTC TGC CCC CCG CTT TTC

1575      1584      1593      1602      1611      1620
CTC TGT TTG AAT ACT ATA AAT TTT CTG GTT CCC TTT GGA TTT ATG GAT AAG TGT

1629      1638
CCC CCT CCA TGT TCC AGG AAA 3'
```

FIGURE 1E

1	M	V	C	R	L	P	V	S	K	K	T	L	L	C	S	F	Q	V	L	D	E	L	G	R	H	V	L	L	R	K	D	C	G	P	V	D	T	K	V	P	consensus
1	M	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	GI 555662	
41	G	A	V	E	P	K	S	A	F	T	Q	G	S	A	M	I	S	S	L	S	O	N	H	P	D	N	R	N	E	T	F	S	S	X	I	S	L	L	N	E	consensus
2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	GI 555662	
81	D	P	L	S	O	D	L	P	V	K	M	A	S	I	F	K	N	F	V	I	T	Y	N	R	T	Y	E	S	K	E	E	A	R	W	R	L	S	V	F	V	consensus
6	E	Y	I	G	F	K	L	E	G	N	V	D	E	K	Y	V	Q	F	K	L	K	Y	R	K	Q	Y	H	E	T	E	D	E	I	-	R	F	N	I	F	K	GI 555662
121	N	N	M	V	R	A	Q	K	I	Q	A	L	D	R	G	T	A	Q	Y	G	V	T	K	F	S	D	L	T	E	E	E	F	R	T	I	Y	L	N	T	L	consensus
45	S	N	I	L	K	A	Q	L	Y	Q	V	F	V	R	G	S	A	I	Y	G	V	T	P	Y	S	D	L	T	D	E	F	A	R	T	H	L	T	A	S	GI 555662	
161	L	R	K	E	P	G	N	K	M	K	O	A	K	S	V	G	-	-	-	D	L	A	P	P	E	W	D	W	R	S	K	G	A	V	T	K	V	K	D	Q	consensus
85	-	W	V	V	P	S	S	R	S	N	T	P	T	S	L	G	K	E	V	N	N	I	P	K	N	F	D	W	R	E	K	G	A	V	T	E	V	K	N	Q	GI 555662
198	G	M	C	G	S	C	W	A	F	S	V	T	G	N	V	E	G	Q	W	F	L	N	Q	G	T	L	L	S	L	S	E	Q	E	L	L	D	C	D	K	M	consensus
124	G	M	C	G	S	C	W	A	F	S	T	T	G	N	V	E	S	Q	W	F	R	K	T	G	K	L	L	S	L	S	E	Q	Q	L	V	D	C	D	G	L	GI 555662
238	D	K	A	C	M	G	G	L	P	S	N	A	Y	S	A	I	K	N	L	G	G	L	E	T	E	D	D	Y	S	Y	Q	G	H	M	Q	S	C	N	F	S	consensus
164	D	D	G	C	N	G	G	L	P	S	N	A	Y	E	S	I	I	K	M	G	G	L	M	L	E	D	N	Y	P	Y	D	A	K	N	E	K	C	H	L	K	GI 555662
278	A	E	K	A	K	V	I	N	D	S	V	E	L	S	Q	N	E	Q	K	L	A	A	W	L	A	E	R	G	P	I	S	V	A	I	N	A	F	G	M	consensus	
204	T	D	G	V	A	V	Y	I	N	S	S	V	N	L	T	Q	D	E	T	E	L	A	A	W	L	Y	H	N	S	T	I	S	V	G	M	N	A	L	L	L	GI 555662

FIGURE 2A

318	Q	F	Y	R	H	G	I	S	R	P	L	C	S	P	W	L	I	D	H	A	V	L	L	V	G	Y	G	-	N	R	S	D	V	P	F	W	A	consensus		
244	Q	F	Y	Q	H	G	I	S	H	P	W	I	F	C	S	K	Y	L	L	D	H	A	V	L	L	V	G	Y	G	V	S	E	K	N	E	P	F	W	I	GI 555662
357	I	K	N	S	W	G	T	D	W	G	E	K	G	Y	Y	L	H	R	G	S	G	A	C	G	V	N	T	M	A	S	S	A	V	V	D	consensus				
284	V	K	N	S	W	G	V	E	W	G	E	N	G	Y	F	R	M	Y	R	G	D	G	S	C	G	I	N	T	V	A	T	S	A	M	I	Y	GI 555662			

FIGURE 2B




FIGURE 3

9/17




FIGURE 4

FIGURE 5A

11/17

387	396	405	414	423	432
TTT GAT CCC AAA GCC TCT AGC TCC TTT CAG GCC AAT GGG ACC AAG TTT GCC ATT					
F D P K A S S S F Q A N G T K F A I					
441	450	459	468	477	486
CAA TAT GGA ACT GGG CGG GTA GAT GAT GGA ATC CTG AGC GAG GAC AAG CTG ACT ATT					
Q Y G T G R V D G I L S E D K L T I					
495	504	513	522	531	540
GGT GGA ATC AAG GGT GCA TCA GTG ATT TTC GGG GAG GCT CTC TGG GAG CCC AGC					
G G I K G A S V I F G E A L W E P S					
549	558	567	576	585	594
CTG GTC TTC GCT TTT GCC CAT TTT GAT GGG ATA TTG GGC CTC GGT TTT CCC ATT					
L V F A F A H F D G I L G L G F P I					
603	612	621	630	639	648
CTG TCT GTG GAA GGA GTT CGG CCC CCG ATG GAT GTA CTG GTG GAG CAG GGG CTA					
L S V E G V R P P P M D V L V E Q G L					
657	666	675	684	693	702
TTG GAT AAG CCT GTC TTC TCC TTT TAC CTC AAC AGG GAC CCT GAA GAG CCT GAT					
L D K P P V F S F Y L N R D P E E P D					
711	720	729	738	747	756
GGA GGA GAG CTG GTC CTG GGG GGC TCG GAC CCG GCA CAC TAC ATC CCA CCC CTC					
G G E L V L G G G S D P A H Y I P P L					

FIGURE 5B

12/17

765	ACC TTC GTG CCA GTC ACG GTC CCC GCC TAC TGG CAG ATC CAC ATG GAG CGT GTG	774	783	792	801	810
	T F V P V T V P A Y W Q I H M E R V					
819	AAG GTG GGC CCA GGG CTG ACT CTC TGT GGC AAG GGC TGT GCT GCC ATC CTG GAT	828	837	846	855	864
	K V G P G L T L C A K G C A A I L D					
873	ACG GGC ACG TCC CTC ATC ACA GGA CCC ACT GAG GAG ATC CGG GCC CTG CAT GCA	882	891	900	909	918
	T G T S L I T G P T E I R A L H A					
927	GCC ATT GGG GGA ATC CCC TTG CTG GCT GGG GAG TAC ATC ATC CTG TGC TCG GAA	936	945	954	963	972
	A I G G I P L L A G E Y I I L C S E					
981	ATC CCA AAG CTC CCC GCA GTC TCC TTC CTT CTT GGG GGC GTC TGG TTT AAC CTC	990	999	1008	1017	1026
	I P K L P A V S F L L G G V W F N L					
1035	ACG GCC CAT GAT TAC GTC ATC CAG ACT ACT ACT CGA AAT GAC GTC TTC TTG GGG ACG	1044	1053	1062	1071	1080
	T A H D Y V I Q T T R N D V F L G T					
1089	TAT GTG GCC GTC TTC GAC CGC GGG GAC ATG AAG AGC AGC GCC CGG GTG GGC CTG	1098	1107	1116	1125	1134
	Y V A V F D R G D M K S S A R V G L					

FIGURE 5C

13/17

1143 GCG CGC GCT CGC ACT CGC GGA GCG GAC CTC GGA TGG GGA GAG ACT GCG CAG GCG 1188
A R A R T R G A D L G W G E T A Q A
1152 1161 1170 1179
1197 1206 1215 1224 1233 1242
CAG TTC CCC GGG TGA CGC CCA AGT GAA GCG CAT GCG CAG CGG GTG GTC GCG GAG
Q F P G *
1251 1260 1269 1278 1287 1296
GTC CTG CTA CCC AGT AAA AAT CCA CTA TTT CCA TTG AAA AAA AAA AAA AAA AAA

AA 3'

FIGURE 5D

1 M S P P P L L Q P L L L L P L L N V E P S G A T L I R I P L H R V Q P G R R I clone312099
 1 M Q P S S L - - - L P L A L C L L A - - A P A S A L V R I P L H K F T S I R R T GI 181180
 41 L N L L R G W - - - - R E P A E L P K L G A P S P G D K P I F V P L S N Y R clone312099
 36 M S E V G G S V E D L I A K G P V S K Y S Q A V P A V T E G P I P E V L K N Y M GI 181180
 75 D V Q Y F G E I G L G T P P Q N F T V A F D T G S S N L W V P S R R C H F F S V clone312099
 76 D A Q Y Y G E I G I G T P P Q C F T V F D T G S S N L W V P S I H C K L L D I GI 181180
 115 P C W L H H R F D P K A S S F O A N G T K F A I O Y G T G R V D G I L S E D K clone312099
 116 A C W I H H K Y N S D K S S T Y V K N G T S F D I H Y G S G S L S G Y L S Q D T GI 181180
 155 L T I - - - - - G G I K G A S V I F G E A L W E P S L V F A F A H F clone312099
 156 V S V P C Q S A S S A S A L G G V K V E R Q V F G E A T K Q P G I T F I A A K F GI 181180
 184 D G I L G L G F P I L S V E G V R P P M D V L V E Q G L L D K P V F S F Y L N R clone312099
 196 D G I L G M A Y P R I S V N N V L E V F D N L M Q Q K L V D Q N I F S F Y L S R GI 181180
 224 D P E E P D G G E L V L G G S D P A H Y I P P L T F V P V T V P A Y W Q I H M E clone312099
 236 D P D A Q P G G E L M L G G T D S K Y Y K G S L S Y L N V T R K A Y W Q V H L D GI 181180
 264 R V K V G P G L T L C A K G C A A I L D T G T S L I T G P T E E I R A L H A A I clone312099
 276 Q V E V A S G L T L C K E G C E A I V D T G T S L M V G P V D E V R E L Q K A I GI 181180

FIGURE 6A

304	GGIPLLAG	EYII	LCSEI	PKLP	AVSF	LLGG	VWFN	LT	AH	DYV	clone312099								
316	GA	VP	LI	QGEY	MI	PC	EKVS	TL	PA	IT	KLGG	KGYK	LS	PE	DYT	GI 181180			
344	IQ	TRN																	
356	LKVS	QAGK	TL	CL	SG	FM	GM	DI	PP	PS	GP	LW	IL	GD	VF	LG	TYVA	clone312099	
																	GR	TYT	GI 181180
359	VFDR	GD	MK	SS	AR	VGL	AR	AR	TR	GA	DL	GW	GE	TA	QA	QF	PG	clone312099	
396	VFDR	DN	-----			N	R	V	CF	AE	A	-----			A	R	L	GI 181180	

FIGURE 6B




FIGURE 7




FIGURE 8

INTERNATIONAL SEARCH REPORT

Internal Application No
PCT/US 97/17175

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N9/64 C12N15/09 C12N15/57 C07K16/40 C07K16/18
A61K38/48 A61K39/395

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	P. FAUST ET AL.: "Cloning and sequence analysis of cDNA for human cathepsin D" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 82, August 1985, pages 4910-4914, XP002052494 whole document	1-28
A	<p style="text-align: center;">---</p> <p>ALENKA PARIS ET AL.: "Molecular cloning and sequence analysis of human preprocathepsin C" FEBS LETTERS, vol. 369, 1995, pages 326-330, XP002011637 whole document, esp. Fig. 3</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	1-28

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

30 January 1998

Date of mailing of the international search report

13. 02. 98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kronester-Frei, A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/17175

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	K. MORI ET AL.: "Molecular cloning of a novel mouse aspartic protease-like protein that is expressed abundantly in the kidney" FEBS LETTERS, vol. 401, 20 January 1997, pages 218-222, XP002052496 whole document	1-28
A	--- WO 95 10630 A (YEDA RES & DEV ;RYCUS AVIGAIL (IL); KIMCHI ADI (IL)) 20 April 1995 whole document, esp. claim 9, Table 1, Fig. 15 -----	1-28

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/17175

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9510630 A	20-04-95	CA 2174136 A	20-04-95
		EP 0804613 A	05-11-97
		JP 9505992 T	17-06-97
